

## **CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge International Advanced Subsidiary and Advanced Level

### **MARK SCHEME for the March 2016 series**

#### **9701 CHEMISTRY**

**9701/52**

Paper 5 (Planning, Analysis and Evaluation),  
maximum raw mark 30

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the March 2016 series for most Cambridge IGCSE® and Cambridge International A and AS Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – March 2016	9701	52

question	expected answer	mark
<b>1 (a)</b>	<p><b>M1</b> (apparatus mark) volumetric flask in range 25–250 cm<sup>3</sup>;</p> <p><b>M2</b> mol propanone = <math>1.00 \times (\text{flask volume} / 1000)</math>; e.g. mol of propanone = <math>1.00 \times 25 / 1000 = 0.025</math> mol</p> <p><b>M3</b> <b>M2</b> <math>\times 58.0</math>; e.g. <math>0.025 \times 58.0 = 1.45</math> g</p>	[3]
<b>(b) (i)</b>	<b>B</b> must be added before first or second reactant	[1]
<b>(ii)</b>	the reactants are <b>A</b> and <b>C</b> so one of these must be mixed last; <b>or</b> the reaction must not start before all three substances are present;	[1]
<b>(c) (i)</b>	(10 cm <sup>3</sup> ) pipette	[1]
<b>(ii)</b>	<p><b>M1</b> NaHCO<sub>3</sub> will effervesce so when effervescence finishes it shows that all H<sup>+</sup> ions have been removed;</p> <p><b>M2</b> NaOH will react with I<sub>2</sub>/CH<sub>3</sub>COCH<sub>3</sub>/reactants;</p>	[2]
<b>(d) (i)</b>	<p><b>M1</b> mol I<sub>2</sub> = <math>(10 / 100) \times 0.200 \times (50 / 1000) = 1.00 \times 10^{-3}</math> mol;</p> <p><b>M2</b> mol S<sub>2</sub>O<sub>3</sub><sup>2-</sup> = <math>2 \times 1.00 \times 10^{-3} = 2.00 \times 10^{-3}</math> mol;</p> <p><b>M3</b> volume 0.100 mol dm<sup>-3</sup> S<sub>2</sub>O<sub>3</sub><sup>2-</sup> = <math>(1000 \times 2.00 \times 10^{-3}) / 0.100 = 20.0</math> cm<sup>3</sup>;</p>	[3]
<b>(ii)</b>	indicator = starch; colour change = blue-black to colourless;	[2]
<b>(e)</b>	time <b>and</b> units of s; volume of thiosulfate <b>and</b> units of cm <sup>3</sup> ;	[2]
<b>(f)</b>	temperature;	[1]
<b>(g) (i)</b>	<p><b>M1</b> (labels) x-axis = time y-axis = concentration of iodine</p> <p><b>M2</b> curved line decreasing from left to right starting from x = 0</p>	[2]

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge International AS/A Level – March 2016	9701	52

question	expected answer	mark																											
(ii)	idea of constant half-life: determine at least two half-lives from the graph and ensure that they are the same; <b>or</b> half-lives determined from the graph should be constant; <b>or</b> determine the gradient (rate) at different points on the graph and plot rate v. concentration to determine if the plot is linear and goes through the origin;	[1]																											
(h)	(incorrect and) half-life will still be constant; <b>or</b> temperature has no effect upon order (of reaction);	[1]																											
2 (a)	<p><b>M1</b> column amount of ethanol burned correctly completed  <b>M2</b> column energy transferred to the water correctly completed</p> <table border="1"> <thead> <tr> <th>experiment number</th><th>amount of ethanol burned / mol</th><th>energy transferred to the water / kJ</th></tr> </thead> <tbody> <tr><td>1</td><td>0.00850</td><td>3.26</td></tr> <tr><td>2</td><td>0.0106</td><td>3.95</td></tr> <tr><td>3</td><td>0.0110</td><td>4.10</td></tr> <tr><td>4</td><td>0.0122</td><td>4.50</td></tr> <tr><td>5</td><td>0.0158</td><td>5.62</td></tr> <tr><td>6</td><td>0.0130</td><td>5.20</td></tr> <tr><td>7</td><td>0.00891</td><td>3.39</td></tr> <tr><td>8</td><td>0.0148</td><td>5.30</td></tr> </tbody> </table>	experiment number	amount of ethanol burned / mol	energy transferred to the water / kJ	1	0.00850	3.26	2	0.0106	3.95	3	0.0110	4.10	4	0.0122	4.50	5	0.0158	5.62	6	0.0130	5.20	7	0.00891	3.39	8	0.0148	5.30	[2]
experiment number	amount of ethanol burned / mol	energy transferred to the water / kJ																											
1	0.00850	3.26																											
2	0.0106	3.95																											
3	0.0110	4.10																											
4	0.0122	4.50																											
5	0.0158	5.62																											
6	0.0130	5.20																											
7	0.00891	3.39																											
8	0.0148	5.30																											
(b)	<b>M1</b> at least eight correctly plotted points; <b>M2</b> correct straight line;	[2]																											
(c)	experiment 6;	[1]																											
(d)	<b>M1</b> co-ordinates, e.g. (0.0106, 3.95) and (0.0158, 5.62); <b>M2</b> gradient correctly calculated from points, e.g. $321 \text{ (kJ mol}^{-1}\text{)}$ ;	[2]																											
(e)	because the reaction is exothermic;	[1]																											
(f) (i)	$((2 \times 0.0005) / 0.391) \times 100 = 0.256\%$ <b>and</b> $(0.05 / 40.0) \times 100 = 0.125\%$ ;	[1]																											

<b>Page 4</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>Cambridge International AS/A Level – March 2016</b>	<b>9701</b>	<b>52</b>

<b>question</b>	<b>expected answer</b>	<b>mark</b>
<b>(ii)</b>	(total) errors in weighing do not account for the (large) error in enthalpy change determined; <b>or</b> heat loss (is more significant);	[1]