



# **Cambridge International Examinations**

Cambridge International Advanced Subsidiary and Advanced Level

CANDIDATE NAME								
CENTRE NUMBER						CANDIDATE NUMBER		
CHEMISTRY								9701/33
Paper 3 Advan	nced Pra	ctical Skil	ls 1				February/I	March 2017
								2 hours
Candidates ans	swer on t	the Quest	tion Pap	er.				
Additional Mate	erials:	As liste	d in the	Confider	ntial Instructions			

### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Give details of the practical session and laboratory where appropriate, in the boxes provided.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 10 and 11. A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [ ] at the end of each question or part question.

Session	
Laboratory	

For Examiner's Use				
1				
2				
3				
Total				

This document consists of 12 printed pages.



1 The concentration of hydrogen peroxide may be given in moldm<sup>-3</sup> or as 'volume strength'. You will determine the concentration of hydrogen peroxide in moldm<sup>-3</sup> and in 'volume strength' by a gas collection method.

Hydrogen peroxide decomposes to form water and oxygen. The reaction is much faster in the presence of a catalyst such as manganese(IV) oxide.

$$2H_2O_2(aq) \rightarrow 2H_2O(I) + O_2(g)$$


'Volume strength' is defined as the volume of oxygen in cm³ produced from the decomposition of 1.0 cm³ of hydrogen peroxide at room temperature and pressure. For example, 1.0 cm³ of '100 volume' hydrogen peroxide will produce 100 cm³ of oxygen.

**FA 1** is a solution of hydrogen peroxide,  $H_2O_2$ . **FA 2** is manganese(IV) oxide,  $MnO_2$ .

## (a) Method

## Read the whole method before starting any practical work.

The diagram below may help you in setting up your apparatus.



- Fill the tub with water to a depth of about 5 cm.
- Fill the 250 cm<sup>3</sup> measuring cylinder **completely** with water. Hold a piece of paper towel firmly over the top, invert the measuring cylinder and place it in the water in the tub.
- Remove the paper towel and clamp the inverted measuring cylinder so that the open end
  is in the water just above the base of the tub.
- Rinse the 50 cm³ measuring cylinder with a little **FA 1** then use it to transfer 150 cm³ of **FA 1** into the reaction flask labelled **X**.
- Check that the bung fits tightly in the neck of flask X, clamp flask X and place the end of the delivery tube into the inverted 250 cm<sup>3</sup> measuring cylinder.
- Remove the bung from the neck of the flask. Tip FA 2 into the hydrogen peroxide and replace the bung immediately. Remove the flask from the clamp and swirl it to mix the contents. Swirl the flask occasionally until no more gas is given off. Replace the flask in the clamp.
- Measure and record the final volume of gas in the measuring cylinder in the space below.

#### Keep FA 1 for use in Question 2.

#### Result

1	(h)	Cal	lcul	latio	ns
М		, va	ıvu	ıatıv	,,,,

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Use the information on page 2 to calculate the 'volume strength' of FA 1.

'volume strength' of **FA 1** = .....

(ii) Calculate the number of moles of oxygen collected in the measuring cylinder. [Assume 1 mole of gas occupies 24.0 dm³ under these conditions.]

moles of  $O_2$  = ..... mol

(iii) Using your answer to (ii) calculate the number of moles of hydrogen peroxide in the volume of **FA 1** added to flask **X**.

moles of  $H_2O_2$  = ..... mol

(iv) Calculate the concentration of hydrogen peroxide, **FA 1**, in mol dm<sup>-3</sup>.

concentration of  $H_2O_2$ , **FA 1** = ..... mol dm<sup>-3</sup> [4]

(c)	(i)	A source of error in this experiment is that some oxygen escapes before the bung can be inserted.
		Suggest a change to the practical procedure given in <b>(a)</b> to reduce this source of error. You may draw a diagram as part of your answer.
	(ii)	The error in reading a $50\text{cm}^3$ measuring cylinder is $\pm 0.5\text{cm}^3$ .
		Calculate the maximum percentage error in the volume of hydrogen peroxide added to flask ${\bf X}$ in ${\bf (a)}$ .
		maximum percentage error in volume of $H_2O_2$ = %
(	(iii)	Explain why the presence of 20 cm³ of air in the 250 cm³ measuring cylinder before the start of the experiment would decrease the accuracy of the results obtained in (a).
		[4]
(d)		ou repeated the method described using half the mass of <b>FA 2</b> , what volume of gas would expect to collect? Explain your answer.

2 You will carry out a second experiment to determine the concentration of hydrogen peroxide, **FA 1**, in mol dm<sup>-3</sup>, by titration with acidified aqueous potassium manganate(VII). The equation for the reaction is given below.

$$2MnO_4^{-}(aq) + 5H_2O_2(aq) + 6H^{+}(aq) \rightarrow 2Mn^{2+}(aq) + 8H_2O(I) + 5O_2(g)$$

**FA 1** is a solution of hydrogen peroxide, H<sub>2</sub>O<sub>2</sub>.

**FA 3** is  $0.0300 \, \text{mol dm}^{-3}$  potassium manganate(VII), KMnO<sub>4</sub>.

FA 4 is dilute sulfuric acid.

# (a) Method

- Fill the burette with **FA 3**.
- Pipette 25.0 cm<sup>3</sup> of **FA 1** into a conical flask.
- Use the 25 cm³ measuring cylinder to add approximately 20 cm³ of FA 4 to the conical flask.
- Perform a rough titration and record your burette readings in the space below.

The rough titre is cr	The rough	titre	is		cm
-----------------------	-----------	-------	----	--	----

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record, in a suitable form below, all of your burette readings and the volume of FA 3
  added in each accurate titration.

I	
II	
III	
IV	
V	
VI	
VII	

[7]

**(b)** From your accurate titration results, obtain a suitable value for the volume of **FA 3** to be used in your calculations. Show clearly how you obtained this value.

25.0 cm<sup>3</sup> of **FA 1** required ...... cm<sup>3</sup> of **FA 3**. [1]

(	C	)	Ca	lc	ul	af	ti	O	n	S
۱		•	Оũ		u	a		v		9

Show y	our working	and appropri	ate significan	t figures in t	the final ar	nswer to <b>each</b>	step of y	your
calculat	tions							

(i)	Calculate the number of moles of manganate(VII) ions present in the volume of	FA	3
	calculated in <b>(b)</b> .		

moles of MnO <sub>4</sub> <sup>-</sup> =	 mol

(ii) Calculate the number of moles of hydrogen peroxide present in 25.0 cm³ of FA 1.

moles of 
$$H_2O_2$$
 = ..... mol

(iii) Using your answer to (ii) calculate the concentration, in mol dm<sup>-3</sup>, of hydrogen peroxide in FA 1.

concentration of  $H_2O_2$  in **FA 1** = ..... mol dm<sup>-3</sup> [4]

[Total: 12]

# 3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs. **No additional tests for ions present should be attempted.** 

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given. (a) FA 5, FA 6 and FA 7 are solutions, some of which contain ions that are listed on pages 10 and 11.

			observations	
	test	FA 5	FA 6	FA 7
(i)	To a 0.5 cm depth of solution in a boiling tube add aqueous sodium hydroxide, then			
	warm gently.			
	Allow to cool, add a piece of aluminium foil and warm again.			
(ii)	To a 1 cm depth of solution in a test-tube add two or three drops of acidified aqueous potassium manganate(VII). (Do <b>not</b> use <b>FA 3</b> .)			
	If no reaction occurs, pour the mixture into a boiling tube and warm gently.			
(iii)	To a 1 cm depth of solution in a test-tube add a 2 cm depth of '10 volume' hydrogen peroxide and leave to stand. (Do <b>not</b> use <b>FA 1</b> .)			
(iv)	To a 1 cm depth of solution in a test-tube add a 1 cm depth of dilute hydrochloric acid, then			
	add a 1 cm depth of aqueous barium chloride or aqueous barium nitrate.			

(b) (i) Identify as many ions present in **FA 5**, **FA 6** and **FA 7** as possible from your observations. If an ion cannot be identified from the tests, write 'unknown' in the space.

	cation(s)	anion(s)
FA 5		
FA 6		
FA 7		

(ii) Describe another test you could carry out to confirm the identity of a cation you have identified in (i). Record the reagent(s) and expected observation(s) in the space below.

Do not carry out this test.

(iii)	Write an ionic equation for the reaction that would occur in (ii). Include state symbols	
		 [6]

[Total:17]

# **Qualitative Analysis Notes**

# 1 Reactions of aqueous cations

	reaction with									
ion	NaOH(aq)	NH <sub>3</sub> (aq)								
aluminium, Al³+(aq)	white ppt. soluble in excess	white ppt. insoluble in excess								
ammonium, NH <sub>4</sub> +(aq)	no ppt. ammonia produced on heating	_								
barium, Ba²+(aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.								
calcium, Ca <sup>2+</sup> (aq)	white ppt. with high [Ca <sup>2+</sup> (aq)]	no ppt.								
chromium(III), Cr³+(aq)	grey-green ppt. soluble in excess	grey-green ppt. insoluble in excess								
copper(II), Cu²+(aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution								
iron(II), Fe <sup>2+</sup> (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess								
iron(III), Fe³+(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess								
magnesium, Mg²+(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess								
manganese(II), Mn <sup>2+</sup> (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess								
zinc, Zn²+(aq)	white ppt. soluble in excess	white ppt. soluble in excess								

# 2 Reactions of anions

ion	reaction
carbonate, CO <sub>3</sub> <sup>2-</sup>	CO <sub>2</sub> liberated by dilute acids
chloride, C <i>l</i> <sup>-</sup> (aq)	gives white ppt. with Ag <sup>+</sup> (aq) (soluble in NH <sub>3</sub> (aq))
bromide, Br <sup>-</sup> (aq)	gives cream ppt. with Ag <sup>+</sup> (aq) (partially soluble in NH <sub>3</sub> (aq))
iodide, I-(aq)	gives yellow ppt. with Ag <sup>+</sup> (aq) (insoluble in NH <sub>3</sub> (aq))
nitrate, NO <sub>3</sub> -(aq)	NH <sub>3</sub> liberated on heating with OH <sup>-</sup> (aq) and A <i>l</i> foil
nitrite, NO <sub>2</sub> -(aq)	$NH_3$ liberated on heating with $OH^-(aq)$ and $Al$ foil; NO liberated by dilute acids (colourless $NO \rightarrow$ (pale) brown $NO_2$ in air)
sulfate, SO <sub>4</sub> <sup>2-</sup> (aq)	gives white ppt. with Ba <sup>2+</sup> (aq) (insoluble in excess dilute strong acids)
sulfite, SO <sub>3</sub> <sup>2-</sup> (aq)	gives white ppt. with Ba <sup>2+</sup> (aq) (soluble in excess dilute strong acids)

# 3 Tests for gases

gas	test and test result
ammonia, NH <sub>3</sub>	turns damp red litmus paper blue
carbon dioxide, CO <sub>2</sub>	gives a white ppt. with limewater (ppt. dissolves with excess CO <sub>2</sub> )
chlorine, Cl <sub>2</sub>	bleaches damp litmus paper
hydrogen, H <sub>2</sub>	'pops' with a lighted splint
oxygen, O <sub>2</sub>	relights a glowing splint

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge International Examinations Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cie.org.uk after the live examination series.

#### The Periodic Table of Elements

Group																	
1	2		13 14 15 16 17												18		
	Key  1 H hydrogen 1.0												2 He helium 4.0				
3	4			atomic numbe				,				5	6	7	8	9	10
Li	Be		ato	mic sym	bol							В	С	N	0	F	Ne
lithium 6.9	beryllium 9.0		rela	name ative atomic m	ass							boron 10.8	carbon 12.0	nitrogen 14.0	oxygen 16.0	fluorine 19.0	neon 20.2
11	12					l						13	14	15	16	17	18
Na	Mg											Αl	Si	Р	S	Cl	Ar
sodium 23.0	magnesium 24.3	3	4	5	6	7	8	9	10	11	12	aluminium 27.0	silicon 28.1	phosphorus 31.0	sulfur 32.1	chlorine 35.5	argon 39.9
19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
potassium 39.1	calcium 40.1	scandium 45.0	titanium 47.9	vanadium 50.9	chromium 52.0	manganese 54.9	iron 55.8	cobalt 58.9	nickel 58.7	copper 63.5	zinc 65.4	gallium 69.7	germanium 72.6	arsenic 74.9	selenium 79.0	bromine 79.9	krypton 83.8
37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
Rb	Sr	Y	Zr	Nb	Мо	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
rubidium 85.5	strontium 87.6	yttrium 88.9	zirconium 91.2	niobium 92.9	molybdenum 95.9	technetium -	ruthenium 101.1	rhodium 102.9	palladium 106.4	silver 107.9	cadmium 112.4	indium 114.8	tin 118.7	antimony 121.8	tellurium 127.6	iodine 126.9	xenon 131.3
55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
Cs	Ва	lanthanoids	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	T <i>l</i>	Pb	Bi	Po	At	Rn
caesium 132.9	barium 137.3		hafnium 178.5	tantalum 180.9	tungsten 183.8	rhenium 186.2	osmium 190.2	iridium 192.2	platinum 195.1	gold 197.0	mercury 200.6	thallium 204.4	lead 207.2	bismuth 209.0	polonium —	astatine -	radon —
87	88	89–103	104	105	106	107	108	109	110	111	112		114		116		
Fr	Ra	actinoids	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn		F1		Lv		
francium	radium		rutherfordium	dubnium	seaborgium	bohrium	hassium	meitnerium	darmstadtium	roentgenium	copernicium		flerovium		livermorium		
_	_		_	_	_	_	_	_	_	_	_		_		_		

lanthanoids	
actinoids	

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Но	Er	Tm	Yb	Lu
lanthanum	cerium	praseodymium	neodymium	promethium	samarium	europium	gadolinium	terbium	dysprosium	holmium	erbium	thulium	ytterbium	lutetium
138.9	140.1	140.9	144.4	-	150.4	152.0	157.3	158.9	162.5	164.9	167.3	168.9	173.1	175.0
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr
actinium	thorium	protactinium	uranium	neptunium	plutonium	americium	curium	berkelium	californium	einsteinium	fermium	mendelevium	nobelium	lawrencium
_	232.0	231.0	238.0	_	_	_	_	-	_	_	_	-	_	_