Cambridge International AS & A Level

Cambridge International Advanced Subsidiary and Advanced Level

	CANDIDATE NAME								
	CENTRE CANDI NUMBER NUMB	DATE ER							
* 6 2	CHEMISTRY	9701/32							
7 2 2 7	Paper 3 Advanced Practical Skills 2	May/June 2016 2 hours							
	Candidates answer on the Question Paper.								
2 4 2 3	Additional Materials: As listed in the Confidential Instructions								
* 🚃	READ THESE INSTRUCTIONS FIRST								
	Write your Centre number, candidate number and name on all the work you had Give details of the practical session and laboratory where appropriate, in the b Write in dark blue or black pen. You may use an HB pencil for any diagrams or graphs. Do not use staples, paper clips, glue or correction fluid. DO NOT WRITE IN ANY BARCODES.	and in. oxes provided.							
	Answer all questions. Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units. Use of a Data Booklet is unnecessary.								
	Qualitative Analysis Notes are printed on pages 10 and 11. A copy of the Periodic Table is printed on page 12.								

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or

Laboratory							
For Exam	iner's Use						
1							

1	
2	
3	
Total	

This document consists of **12** printed pages.



part question.

1 In **Questions 1** and **2** you will determine the relative atomic mass, *A*_r, of magnesium by two different methods.

In the first method you will collect and measure the volume of gas given off in the reaction between a known mass of magnesium and a known amount of dilute sulfuric acid. The acid will be in excess.

 $Mg(s) + H_2SO_4(aq) \rightarrow MgSO_4(aq) + H_2(g)$

FB 1 is a strip of magnesium ribbon, Mg. **FB 2** is 1.00 mol dm⁻³ sulfuric acid, H₂SO₄.

(a) Method

Read through the whole method before starting any practical work.

The diagrams below may help you in setting up your apparatus.



- Fill the tub with water to a depth of about 5 cm.
- Fill the 250 cm³ measuring cylinder **completely** with water. Hold a piece of paper towel firmly over the top, invert the measuring cylinder and place it in the water in the tub.
- Remove the paper towel and clamp the inverted measuring cylinder so the open end is in the water just above the base of the tub.
- Pipette 25.0 cm³ of **FB 2** into the reaction flask labelled **X**.
- Check that the bung fits tightly in the neck of flask **X**, clamp flask **X**, and place the end of the delivery tube into the inverted 250 cm³ measuring cylinder.
- Weigh the magnesium ribbon, **FB 1**, and record the mass in the space below.
- Remove the bung from the neck of the flask. Add the magnesium ribbon, **FB 1**, into the acid in the flask and replace the bung **immediately**. Remove the flask from the clamp and swirl it to mix the contents. Swirl the flask occasionally until no more gas is evolved. Replace the flask in the clamp.
- When no more gas is collected, measure and record the final volume of gas in the measuring cylinder in the space below.

KEEP THE CONTENTS OF THE CONICAL FLASK X FOR USE IN QUESTION 2.

Results

(b) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Calculate the number of moles of gas collected in the measuring cylinder. [Assume that 1 mole of gas occupies 24.0 dm³ under these conditions.]

moles of gas = mol

(ii) Use your answer to (i) and the mass of magnesium, **FB 1**, recorded in (a) to calculate the relative atomic mass, *A*_r, of magnesium.

A_r of magnesium =

[3]

[Total: 5]

2 You will determine the amount of sulfuric acid remaining in flask **X** after the reaction with magnesium in **Question 1**. You will do this by titration with sodium hydroxide of known concentration.

 $2NaOH(aq) + H_2SO_4(aq) \rightarrow Na_2SO_4(aq) + 2H_2O(I)$

FB 3 is 0.150 mol dm⁻³ sodium hydroxide, NaOH. bromophenol blue indicator

(a) Method

- Transfer all the contents of flask **X** from **Question 1** into the 250 cm³ volumetric flask.
- Rinse flask **X** with distilled water and add the washings to the volumetric flask. Add distilled water up to the mark. Stopper the volumetric flask and mix the contents thoroughly.
- Label this solution **FB 4**.
- Rinse the pipette and use it to transfer 25.0 cm³ of **FB 4** into the conical flask.
- Add about 10 drops of bromophenol blue to the conical flask.
- Fill the burette with **FB 3**.
- Perform a rough titration and record your burette readings in the space below. The end point is reached when the solution becomes a permanent blue-violet colour.

The rough titre is cm³.

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Record, in a suitable form below, all of your burette readings and the volume of FB 3 added in each accurate titration.
- Make certain any recorded results show the precision of your practical work.



[7]

(b) From your accurate titration results, obtain a suitable value for the volume of **FB 3** to be used in your calculations. Show clearly how you obtained this value.

25.0 cm³ of **FB 4** required cm³ of **FB 3**. [1]

(c) Calculations

Show your working and appropriate significant figures in the final answer to **each** step of your calculations.

(i) Calculate the number of moles of sodium hydroxide, NaOH, in the volume of **FB 3** you calculated in (b).

moles of NaOH = mol

(ii) Use your answer to (i) to calculate the number of moles of sulfuric acid present in the 25.0 cm³ of **FB 4** pipetted in (a).

moles of H_2SO_4 = mol

(iii) Use your answer to (ii) to calculate the number of moles of sulfuric acid, H₂SO₄, remaining in flask **X** after the reaction in **1(a)**.

moles of H_2SO_4 remaining from **1(a)** = mol

(iv) Use the relevant information on page 2 to calculate the number of moles of sulfuric acid, H_2SO_4 , pipetted into reaction flask **X** in **1(a)**.

moles of H_2SO_4 pipetted into flask **X** = mol

(v) Use your answers to (iii) and (iv) to calculate the number of moles of sulfuric acid which reacted with the magnesium in flask X.

moles of H_2SO_4 which reacted in flask **X** = mol

(vi) Use your answer to (v) and the mass of magnesium used in **1(a)** to calculate the relative atomic mass, *A*_r, of magnesium.

A_r of magnesium =[6]

(d) (i) A student, who carried out the experiments in **Questions 1** and **2** correctly, calculated the A_r of magnesium as shown in the table.

	Question 1	Question 2				
A _r Mg	20.8	22.5				

Use the A_r of magnesium given in the Periodic Table on page 12 to deduce which practical procedure is less accurate. Identify one source of inaccuracy and explain **one** change the student could make in order to improve the accuracy.

	is less accurate
	source of inaccuracy
	improvement
(ii)	Use the A_r of magnesium given in the Periodic Table to calculate the percentage error in the student's value from Question 1 .
	[3]

[Total: 17]

3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs. Marks are **not** given for chemical equations. **No additional tests for ions present should be attempted.**

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

(a) (i) **FB 5** is a solid element and **FB 6** is a solid compound containing one cation and one anion. Carry out the following tests and record your observations.

test	observations
Place a small spatula measure of FB 5 in a boiling tube, add a 1 cm depth of dilute hydrochloric acid and warm the contents of the tube gently.	
Place small spatula measures of FB 5 and FB 6 in a single boiling tube. Use a test-tube holder to hold the tube. Add a 2 cm depth of aqueous sodium hydroxide. CARE	
Place a spatula measure of FB 6 in a test-tube. Add a 3 cm depth of distilled water to form a solution for the following two tests.	
To a 1 cm depth of aqueous FB 6 in a test-tube add a 1 cm depth of dilute hydrochloric acid.	
To a 1 cm depth of aqueous FB 6 in a test-tube add aqueous sodium hydroxide.	

(ii) Suggest the identities of **FB 5** and **FB 6** from your observations. Refer to the relevant observations in your answers.

FB 5:	
reason(s)	
FB 6: cation	reason(s)
anion	reason(s)
	[9]

- (b) (i) FB 7 and FB 8 are solutions which contain different anions. These may be carbonate, halide, sulfate or sulfite. You are to devise tests to identify the two anions present. Record in a suitable table below:
 - the reagent(s) you use for each test,
 - the observations you make on carrying out the test,
 - the conclusion you make from the result of the test.

(ii) The cation in either **FB 7** or **FB 8** is a transition metal ion. Carry out the following test to identify this cation and record your observations.

toot	observations							
lest	FB 7	FB 8						
To a 1 cm depth of solution in a test-tube add aqueous sodium hydroxide.								

FB contains the transition metal ion

[9]

[Total: 18]

Qualitative Analysis Notes

Key: [ppt. = precipitate]

1 Reactions of aqueous cations

ion	reaction	on with
юп	NaOH(aq)	NH ₃ (aq)
aluminium, A <i>l</i> ³⁺(aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH₄⁺(aq)	no ppt. ammonia produced on heating	_
barium, Ba²⁺(aq)	faint white ppt. is nearly always observed unless reagents are pure	no ppt.
calcium, Ca²⁺(aq)	white ppt. with high [Ca ²⁺ (aq)]	no ppt.
chromium(III), Cr³⁺(aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu²+(aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe²+(aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg²⁺(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn²⁺(aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn²⁺(aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

ion	reaction
carbonate, CO ₃ ²-	CO ₂ liberated by dilute acids
chloride, C <i>l⁻</i> (aq)	gives white ppt. with Ag $^{+}(aq)$ (soluble in NH $_{3}(aq))$
bromide, Br⁻(aq)	gives cream ppt. with Ag ⁺ (aq) (partially soluble in $NH_3(aq)$)
iodide, I⁻(aq)	gives yellow ppt. with Ag $^{*}(aq)$ (insoluble in NH $_{3}(aq)$)
nitrate, NO ₃ ⁻(aq)	NH_3 liberated on heating with $OH^-(aq)$ and Al foil
nitrite, NO₂⁻(aq)	NH_3 liberated on heating with OH ⁻ (aq) and A <i>l</i> foil; NO liberated by dilute acids (colourless NO \rightarrow (pale) brown NO ₂ in air)
sulfate, SO ₄ ²-(aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids)
sulfite, SO ₃ ²-(aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids)

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater (ppt. dissolves with excess CO_2)
chlorine, Cl_2	bleaches damp litmus paper
hydrogen, H ₂	"pops" with a lighted splint
oxygen, O ₂	relights a glowing splint

Group 1 2	The Periodic Table of Elements																	
1 2 13 14 15 16 17 1 3 4 15 16 17 1 1 1 16 17 1 3 4 15 16 17 1 1 14 15 16 17 1 3 4 15 16 17 1 1 16 17 1 14 15 16 17 1 14 15 16 17 1 14 15 16 17 1 14 15 16 17 1 14 15 16 17 1 16 17 1 16 17 1 14 15 16 17 1 16 17 1 16 17 1 16 17 1 16 16 17 16 17 16 17 16 17 16 17 16 16 16	Group																	
interm Key 3 4 atomic number atomic symbol atomic number atomic number <td< td=""><td>1</td><td>2</td><td colspan="11">13 14 15 16 17 7</td><td>18</td></td<>	1	2	13 14 15 16 17 7											18				
3 4 atomic number atomic number 11 12 11 12 11 12 11 12 13 14 15 16 17 1 10 11 12 13 14 15 16 17 1 11 12 13 14 15 16 17 1 120 24.3 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 1 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 44 35 3 34 35 3 34 35 44 35 45 46 47 46 49 30 15 22 35 55 55.9 55.9 55.9 55.9 55.9 55.9 55.9 75.9	Key 1 H hydrogen 1.0												2 He helium 4.0					
Li Be many relative atomic symbol model Limbur method 11 Limbur method 12.0 Be many relative atomic mass C N O F N N 11 12 Na Mg - N - <td>3</td> <td>4</td> <td></td> <td></td> <td>atomic numbe</td> <td>r</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td>5</td> <td>6</td> <td>7</td> <td>8</td> <td>9</td> <td>10</td>	3	4			atomic numbe	r							5	6	7	8	9	10
Itelan Desptisus relative adorim respective address relative adorim relative adorim relative address relative address <threlative address relative address<!--</td--><td>Li</td><td>Be</td><td></td><td>ato</td><td>omic sym</td><td>bol</td><td></td><td></td><td></td><td></td><td></td><td></td><td>В</td><td>С</td><td>N</td><td>0</td><td>F</td><td>Ne</td></threlative 	Li	Be		ato	omic sym	bol							В	С	N	0	F	Ne
11 12 13 14 15 16 17 1 Naduum 24.3 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 1 19 20 21 22 23 24 25 26 27 28 29 30 31 32.3 34 35 3 19 20 21 22 23 24 25 26 27 28 29 30 31 32.3 34 35 3 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 5 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 5	lithium 6.9	beryllium 9.0		rel	name ative atomic m	ass							boron 10.8	carbon 12.0	nitrogen 14.0	oxygen 16.0	fluorine 19.0	neon 20.2
Na Mg 23.0 Mg 23.0 No Al Si P S Cl A 23.0 24.3 3 4 5 6 7 8 9 10 11 12 allowand position askin position	11	12					1						13	14	15	16	17	18
solution magnetisming 3 4 5 6 7 8 9 10 11 12 attioning attion attion in an analysis 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 3 gatesian Case Sc Ti V Cr M Fe Co Ni Cu Zn Ga Ge As See Br K 931 40.1 41 42 43 44 46 47 48 49 50 51 52 53 55 56 57-71 78 79.0 77.3 74 75 77 78 79 80 81 82 83 84 85 8 76 67.6 85.7 71 72 73 74 75 76 77 78 79	Na	Mg											Al	Si	Р	S	Cl	Ar
19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 94 35 3 potessium 381 44.0 50.9 27 28 29 30 21 32 33 94 35 3 37 38 39 40 41 42 43 44 45 46 47 48 49 51 52 53 55 56 57.71 72.7 Nb Mo Te I Xz 27.7 28 79 80 77 78 79 80 81 82 83 84 85 8 10404m stortium risclam rodium	sodium 23.0	magnesium 24.3	3	4	5	6	7	8	9	10	11	12	aluminium 27.0	silicon 28.1	phosphorus 31.0	sulfur 32.1	chlorine 35.5	argon 39.9
K Ca Sc Ti V Cr Mn Fe Co Ni Cu Zn Ga Ge As Se Br Write 33.1 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 5 Rb Sr Y Zr Nb Mo Tc Ru Rh Pdd Add	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36
protestion cardium scandium standium	K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 55 Nbb witnitum 85.5 87.6 87.6 91.2 92.9 91.2 73 74 75 76 77 78 79 80 81 82 83 84 85 85 126.9 132.8 127.6 126.9 132.8 127.6 126.9 126.9 132.8 127.8 127.6 126.9 132.8 127.6 126.9 132.8 127.8 127.6 126.9 132.8 127.8 127.6 126.9 132.8 132.8 132.8 127.8 126.9 126.9 126.9 126.9 126.9 126.9 126.9 127.8 127.6 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9 126.9<	potassium 39.1	calcium 40.1	scandium 45.0	titanium 47.9	vanadium 50.9	chromium 52.0	manganese 54.9	iron 55.8	cobalt 58.9	nickel 58.7	copper 63.5	zinc 65.4	gallium 69.7	germanium 72.6	arsenic 74.9	selenium 79.0	bromine 79.9	krypton 83.8
Rb nubicium 85.5 Sr 87.6 Y 88.9 Zr 91.2 Nb 92.9 Mo 99.9 Tc exhibitium 92.9 Ru 99.9 Ru technetium 101.1 Ru 102.8 Rd palladium 102.9 Cd silver 102.4 Indum 112.4 Sn 112.4 Sn 12.6 Sn 12.6 Sn 12.6 Sn 12.6 Sn 12.6 Sn 12.6 Sn 13.7 S	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54
	Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
55 56 57-71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 8 Cs Ba barrium Ianthanoids Hf Ta W Re Os Ir Pt Au Hg Tl Pb Bi Po At astatime radium 132.9 137.3 104 105 106 107 108 109 110 111 112 114 116 astatime radium radium astatime radium 197.0 200.4 207.2 114 116 Image: Neget Act	rubidium 85.5	strontium 87.6	yttrium 88.9	zirconium 91.2	niobium 92.9	molybdenum 95.9	technetium -	ruthenium 101.1	rhodium 102.9	palladium 106.4	silver 107.9	cadmium 112.4	indium 114.8	tin 118.7	antimony 121.8	tellurium 127.6	iodine 126.9	xenon 131.3
Cs Ba Ianthanoids Hf Ta W Re Os Ir Pt Au Hg Tl Pb Bi Po At Restance 132.9 137.3 137.3 104 105 106 107 108 109.2 195.1 197.0 200.6 204.4 209.0 -	55	56	57–71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86
caseJun 132.9 barrum 137.3 hafnium 178.5 tandium 180.9 tungsten 180.9 menium 180.2 indium 190.2 platinum 192.2 polo mercury 197.0 thallium 200.6 lead 204.4 bismuth 207.2 polonium 200.0 astaline 204.4 polonium 201.2 platinum 200.6 lead 204.4 platinum 207.2 polonium 200.0 method 204.4 platinum 201.2 lead 204.4 platinum 201.2 lead 204.4 platinum 201.2 platinum 200.6 platinum 204.4 platinum 207.2 platinum 200.0 platinum 204.4 platinum 201.4 platinum 200.6 platinum 204.4 platinum 201.4 platinum 40.4	Cs	Ba	lanthanoids	Hf	Та	W	Re	Os	Ir	Pt	Au	Hg	Τl	Pb	Bi	Po	At	Rn
Solide Other Other <t< td=""><td>caesium 132.9</td><td>barium 137.3</td><td></td><td>hafnium 178.5</td><td>tantalum 180.9</td><td>tungsten 183 8</td><td>rhenium 186 2</td><td>osmium 190.2</td><td>iridium 192.2</td><td>platinum 195 1</td><td>gold 197.0</td><td>mercury 200.6</td><td>thallium 204 4</td><td>lead 207.2</td><td>bismuth 209.0</td><td>polonium</td><td>astatine</td><td>radon</td></t<>	caesium 132.9	barium 137.3		hafnium 178.5	tantalum 180.9	tungsten 183 8	rhenium 186 2	osmium 190.2	iridium 192.2	platinum 195 1	gold 197.0	mercury 200.6	thallium 204 4	lead 207.2	bismuth 209.0	polonium	astatine	radon
Fr franciumRa radiumactinoidsRf rutherfordiumDb dubniumSg seaborgiumBh bohriumHs hassiumMt melineriumDs darmstadtiumRg rentgeniumCn coperniciumFl theroviumLv livermoriumIanthanoids575859606162636465666768697071IanthanoidsLa lanthanumCe ceriumPr 140.1Nd reserviniumPm reserviniumSm samarium 150.4Eu 150.4Gd 152.0Tb gadolinium 157.3Dy terbiumHo terbium terbiumFr terbium tholinum tholinum tholinum terbiumMd reservinium 162.5Nd terbium terbiumPm terbium terbium terbium tist.9Gd terbium terbium terbium tist.9162.5666768697071 tuu tuu lutuitum tuu lutuitum tuu lutuitum tuu lutuitum trast168.97071 tuu tuu lutuitum tuu lutuitum tuu lutuitum tuu lutuitum trastactinoids575859606162636465666768697071 tuu lutuitum tuu lutuitum tuu lutuitum tuu lutuitum tuu tuu lutuitum tuu lutuitum tuu lutuitum promethium protectinium protectinium tur protectinium tur protectinium tur tur tur protectinium tervinum60666768697071 tuu tuu <b< td=""><td>87</td><td>88</td><td>89–103</td><td>104</td><td>105</td><td>106</td><td>107</td><td>108</td><td>109</td><td>110</td><td>111</td><td>112</td><td>20111</td><td>114</td><td>200.0</td><td>116</td><td></td><td></td></b<>	87	88	89–103	104	105	106	107	108	109	110	111	112	20111	114	200.0	116		
francium radium rutherfordium dubnium seaborgium bohrium hassium meilnerium darmstadtium rcentgenium copernicium flerovium livermorium -	Fr	Ra	actinoids	Rf	Db	Sg	Bh	Hs	Mt	Ds	Rg	Cn		F1		Lv		
Ianthanoids 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 Ianthanoids La Ce Pr Nd Pm Sm Eu Gd Tb Dy Ho Er Tm Yb Lu Ianthanum 138.9 140.1 140.9 144.4 - 150.4 152.0 157.3 158.9 162.5 164.9 167.3 168.9 173.1 175.0 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 actinoids Ac Th Pa U Np Pu Am Cm Bk Cf Es Fm Md No Lr actinoids 231.0 231.0 231.0 231.0 231.0 231.0 231.0 231.0 231.0 231.0 231.0	francium	radium		rutherfordium	dubnium	seaborgium	bohrium	hassium	meitnerium	darmstadtium	roentgenium	copernicium		flerovium		livermorium		
Ianthanoids 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 Ianthanoids Ianthanum Ce lanthanum Pr Nd receium Pm Sm Eu samarium Gd Tb Dy Ho Er Tm Yb Lu 138.9 140.1 140.9 144.4 - 150.4 157.3 158.9 162.5 164.9 167.3 168.9 173.1 175.0 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 actinoids Ac Th Pa U Np Pu Am Cm Bk Cf Es Fm Md No Lr actinium thorium protactinum urenium cultum cultum cultum cultum cultum cultum cultum cultum			<u> </u>					_	_						1			
IanthanoidsLa lanthanumCe ceriumPr prasedymiumNd nedymiumPm nedymiumSm samariumEu europiumGd gddiniumTb terbiumDy dysprosiumHo holmiumEr erbiumTm thuliumYb yterbiumLu lutetium138.9140.1140.9144.4-150.4152.0157.3158.9162.5164.9167.3168.9173.1175.08990919293949596979899100101102103actinoidsAc actiniumTh thoriumPa uraniumU uraniumNp neptuniumPu plutoniumAm americiumCm curiumBk curiumCf califormiumEs celliumFm mendeleviumMd nobeliumNo nobeliumLr lawrencium23102310238.0			57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	
Ianthanum cerium praseodymium neodymium promethium samarium europium gadolinium terbium dysprosium holmium erbium thulium ytterbium tultulum ytterbium lutetlum 138.9 140.1 140.9 144.4 - 150.4 157.3 157.3 158.9 162.5 164.9 167.3 168.9 173.1 175.0 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 Acc Th Pa U Np Pu Am Cm Bk Cf Es Fm Md No Lr actinium thorium protactinium uranium protactinium protactinium gadolinium curium curium curium fertilium fertilium fertilium mendelevium nobelium lawrencium actinium thorium 231.0 231.0 231.0 231.0 231.0 231.0 231.0 231.0 231.0 231.0	lanthanoids		La	Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu	
actinoids8990919293949596979899100101102103Act actinium thorium thorium 233.0Th protactinium 233.0Pa uranium 233.0U reputnium americium americium americium96979899100101102103Bk curium protactinium actinium curiumPa protactinium americium 233.0949596979899100101102103Bk berkelium curiumCf curium curium berkelium curiumBk curium curium curium curium einsteinium fermium fermium fermium fermium fermium curium100101102103Bk curium plutonium curium plutonium curium curium curium plutonium curium 			lanthanum 138.9	cerium 140.1	praseodymium 140.9	neodymium 144.4	promethium -	samarium 150.4	europium 152.0	gadolinium 157.3	terbium 158.9	dysprosium 162.5	holmium 164.9	erbium 167.3	thulium 168.9	ytterbium 173.1	lutetium 175.0	
actinoids Ac Th Pa U Np Pu Am Cm Bk Cf Es Fm Md No Lr actinium thorium 233.0 231.0 231.0 231.0	89 90 91 92 93 94 95 96 97 98 99 100 101 102 10					103												
actinium thorium protactinium uranium neptunium plutonium americium curium berkelium californium einsteinium fermium mendelevium nobelium lawrencium	actinoid	s	Ac	Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr	
			actinium -	thorium 232.0	protactinium 231.0	uranium 238.0	neptunium —	plutonium —	americium -	curium —	berkelium -	californium -	einsteinium -	fermium —	mendelevium -	nobelium —	lawrencium -	