
CHEMISTRY

9701/22

Paper 2 AS Level Structured Questions

May/June 2017

MARK SCHEME

Maximum Mark: 60

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2017 series for most Cambridge IGCSE[®], Cambridge International A and AS Level and Cambridge Pre-U components, and some Cambridge O Level components.

PUBLISHED

Question	Answer						Marks
1(a)	atomic number	nucleon number	number of electrons	number of protons	number of neutrons	symbol	2 1 1
		6		3	3		
						$^{58}_{26}\text{Fe}^{3+}$	
1(b)(i)	EITHER mass of an atom / isotope relative / compared to 1/12 (the mass) of (an atom of) C-12 OR on a scale in which a C-12 (atom / isotope) has (a mass of exactly) 12 (units) OR mass of one mol (of atoms) of an isotope relative / compared to 1/12 (the mass) of 1 mol of C-12 OR on a scale in which one mol C-12 (atom / isotope) has a mass of (exactly) 12 g						2 1 1
1(b)(ii)	$\frac{(10.0129 \times 19.78) + (80.22x)}{100} = 10.8$						1
	x = 10.9941						1
	Total:						6

PUBLISHED

Question	Answer	Marks
2(a)	strong triple bond	1
	non-polar / no dipole	1
2(b)(i)	Any 2 points covered correctly scores 2 marks Any 1 point covered correctly scores 1 mark <ul style="list-style-type: none"> nitrogen (and oxygen) from the air / atmosphere (react): high temperature (of internal combustion engine) / (engine) produces enough OR a lot of heat (energy) : (so) breaks (strong) bond(s) in nitrogen (and oxygen) : 	2
2(b)(ii)	reduction / decomposition of NO _x using a catalyst / catalytic convertor	1
	2NO ₂ + 4CO → 4CO ₂ + N ₂ OR 2NO + 2CO → 2CO ₂ + N ₂	1
2(b)(iii)	(acts as a homogeneous) catalyst OR oxidising agent	1
	SO ₂ + NO ₂ → SO ₃ + NO	1
	NO + ½O ₂ → NO ₂ OR SO ₃ + H ₂ O → H ₂ SO ₄	1
2(b)(iv)	2NO ₂ + H ₂ O → HNO ₂ + HNO ₃ OR 4NO ₂ + 2H ₂ O + O ₂ → 4HNO ₃	1
2(c)	fertiliser / nitrates dissolve in (river water) OR fertiliser / nitrates are washed / leached out / flows into (river water)	1

PUBLISHED



Question	Answer	Marks
	algal bloom / promote algal growth / explosion of plant growth AND EITHER sunlight is blocked out (preventing photosynthesis) / plants can no longer carry out photosynthesis (and die) OR bacteria break down or decay dead organisms / plants / algae	1
	drop in oxygen (concentration)	1
	Total:	13


PUBLISHED

Question	Answer	Marks
3(a)	(+) 103	1
3(b)(i)	general shape of the curve and peak are displaced to right of original and starts at origin	1
	the peak is lower and curve crosses once only finishing above original	1
3(b)(ii)	rate increases AND correct explanation in terms of 'more collisions'	1
	at higher T area above E_a is greater / more molecules with $E \geq E_a$	1
	higher frequency of successful collisions OR more successful collisions per unit time / higher chance of successful collisions per unit time / higher proportion of successful collisions per unit time	1
3(b)(iii)	increases (%) decomposition (of HBr)	1
	(increasing T) shifts equilibrium to the right / in the forward direction / endothermic direction / towards $H_2 + Br_2$	1
	to oppose the change or oppose the increase in temperature OR to absorb (additional) energy / heat OR to decrease the temperature	1
3(b)(iv)	H-I bond strength less than H-Br OR less energy needed to break H-I <i>ora</i>	1
	I (atom) is big(ger) (than Br) OR I (atom) has more shielding (than Br) <i>ora</i>	1
	Br (atom) has greater (%) orbital / outer shell overlap OR attraction (of nucleus in iodine) for shared (pair of) electrons is weak(er) OR attraction (of nucleus in iodine) for bonding pair (or electrons) is weak(er) <i>ora</i>	1

PUBLISHED

Question	Answer	Marks
3(c)(i)	$H_2 = 0.015 \text{ (mol)}$	1
	$HCl = 0.27 \text{ (mol)}$	1
3(c)(ii)	$HCl = 9/10$ AND $x_{H_2} = 1/20$ AND $C_{l_2} = 1/20$ OR $HCl = 0.9(0)$ AND $H_2 = 0.05$ AND $C_{l_2} = 0.05$	1
3(d)(i)	$(K_p =) \frac{p_{H_2} \times p_{Cl_2}}{p_{HCl}^2}$	1
3(d)(ii)	equal number of moles (of gas) on either side (of equation) / (total) pressure cancels	1
3(d)(iii)	4.649×10^{-3}	1
	Total:	18

Question	Answer	Marks
4(a)(i)	(A =) 	1
4(a)(ii)	(A / straight chain) has strong(er) (temporary dipole-) induced dipole (attractions) <i>ora</i>	1
	(because A / straight chain has) bigger (surface) area / more (points of) contact (in unbranched isomer) <i>ora</i> OR (so) more energy required to break the intermolecular forces <i>ora</i>	1
4(a)(iii)	CH ₃ CHCHCH ₃ OR CH ₃ CH=CHCH ₃	1
4(a)(iv)	No rotation / restricted / limited rotation of C=C / (carbon) double bond	1
	One (of the two) methyl groups / one (of the two) H (atoms) is on each C (of C=C)	1
4(a)(v)		1
	arrow from the C=C double bond drawn to the bromine	1
	dipole on Br ₂ in correct orientation AND arrow from the Br-Br bond to the Br ^{δ-}	1
	correct carbocation / bromonium ion from the structure with C=C drawn OR Br ⁻ with lone pair, negative charge AND arrow from lone pair to the carbon atom of intermediate OR using both arrows shown (in alternative diagram)	1
4(a)(vi)	electrons in pi bond induce it (the dipole) OR (high) electron density in pi bond / double bond / C=C repels electrons (away from nearest Br) OR polarised by (high) electron density in pi bond / double bond / C=C	1

Question	Answer	Marks
4(b)(i)	C = (2-)methylpropan-2-ol / $(\text{CH}_3)_3\text{COH}$ / any unambiguous structure	1
	D = (2-)methylpropan-1-ol / $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$ / any unambiguous structure	1
	E = (2-)methylpropanoic acid / $(\text{CH}_3)_2\text{CHCO}_2\text{H}$ / any unambiguous structure	1
	 <p style="text-align: center;"> C D E </p>	
4(b)(ii)	$2\text{C}_4\text{H}_8\text{O}_2 + \text{Na}_2\text{CO}_3 \rightarrow 2\text{C}_4\text{H}_7\text{O}_2\text{Na} + \text{H}_2\text{O} + \text{CO}_2$	1
4(c)(i)	triiodomethane	1
4(c)(ii)	F = $\text{CH}_3\text{CH}_2\text{CH}_2\text{COCH}_3$	1
	G = $\text{C}_2\text{H}_5\text{CH}(\text{CH}_3)\text{CHO}$	1
4(c)(iii)	a (tetrahedral) atom with four different groups / atoms / substituents attached OR a carbon (atom) with four different groups / atoms / substituents attached	1
4(d)(i)	H $\text{C}=\text{O}$ (group / bond) AND $\text{O}-\text{H}$ (group / bond)	1
	I $\text{C}=\text{O}$ (group / bond) AND $\text{C}-\text{H}$ (group / bond)	1

PUBLISHED

Question	Answer	Marks
4(d)(ii)	H = ethanoic acid	1
	I = methyl methanoate	1
	Total:	23