Auv	anced Subsidiary Level and Ac	avanced Level
MATHEMATICS		9709/0 <sup>-</sup>
Paper 1 Pure Ma	thematics 1 (P1)	May/June 200
Additional Materials:	Answer Booklet/Paper Graph paper List of Formulae (MF9)	1 hour 45 minute

## **READ THESE INSTRUCTIONS FIRST**

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet. Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen on both sides of the paper. You may use a soft pencil for any diagrams or graphs. Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all the questions.

Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.

The use of an electronic calculator is expected, where appropriate.

You are reminded of the need for clear presentation in your answers.

The number of marks is given in brackets [] at the end of each question or part question. The total number of marks for this paper is 75.

Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

At the end of the examination, fasten all your work securely together.

## This document consists of **4** printed pages.



2 Solve the equation

$$\sin 2x + 3\cos 2x = 0,$$

[4]


[2]

for  $0^{\circ} \le x \le 180^{\circ}$ .

3 Each year a company gives a grant to a charity. The amount given each year increases by 5% of its value in the preceding year. The grant in 2001 was \$5000. Find

- (ii) the total amount of money given to the charity during the years 2001 to 2011 inclusive. [2]
- 4 The first three terms in the expansion of  $(2 + ax)^n$ , in ascending powers of x, are  $32 40x + bx^2$ . Find the values of the constants *n*, *a* and *b*. [5]
- 5 The curve  $y^2 = 12x$  intersects the line 3y = 4x + 6 at two points. Find the distance between the two points. [6]






In the diagram, *ABC* is a triangle in which AB = 4 cm, BC = 6 cm and angle  $ABC = 150^{\circ}$ . The line *CX* is perpendicular to the line *ABX*.

(i) Find the exact length of *BX* and show that angle 
$$CAB = \tan^{-1}\left(\frac{3}{4+3\sqrt{3}}\right)$$
. [4]


(ii) Show that the exact length of AC is  $\sqrt{(52 + 24\sqrt{3})}$  cm.



3

The diagram shows a circle with centre *O* and radius 8 cm. Points *A* and *B* lie on the circle. The tangents at *A* and *B* meet at the point *T*, and AT = BT = 15 cm.

- (i) Show that angle *AOB* is 2.16 radians, correct to 3 significant figures. [3]
- (ii) Find the perimeter of the shaded region.
- (iii) Find the area of the shaded region.



The diagram shows the roof of a house. The base of the roof, OABC, is rectangular and horizontal with OA = CB = 14 m and OC = AB = 8 m. The top of the roof DE is 5 m above the base and DE = 6 m. The sloping edges OD, CD, AE and BE are all equal in length.

Unit vectors **i** and **j** are parallel to OA and OC respectively and the unit vector **k** is vertically upwards.

- (i) Express the vector  $\overrightarrow{OD}$  in terms of i, j and k, and find its magnitude. [4]
- (ii) Use a scalar product to find angle *DOB*.

7

8

[4]


[2]

[3]

9 A curve is such that  $\frac{dy}{dx} = \frac{4}{\sqrt{6-2x}}$ , and P(1, 8) is a point on the curve.

- (i) The normal to the curve at the point P meets the coordinate axes at Q and at R. Find the coordinates of the mid-point of QR. [5]
- (ii) Find the equation of the curve.

10



The diagram shows the curve  $y = x^3 - 3x^2 - 9x + k$ , where k is a constant. The curve has a minimum point on the x-axis.

- (i) Find the value of k. [4]
- (ii) Find the coordinates of the maximum point of the curve. [1]
- (iii) State the set of values of x for which  $x^3 3x^2 9x + k$  is a decreasing function of x. [1]
- (iv) Find the area of the shaded region.

## **11** Functions f and g are defined by

f:  $x \mapsto k - x$  for  $x \in \mathbb{R}$ , where k is a constant, g:  $x \mapsto \frac{9}{x+2}$  for  $x \in \mathbb{R}$ ,  $x \neq -2$ .

- (i) Find the values of k for which the equation f(x) = g(x) has two equal roots and solve the equation f(x) = g(x) in these cases. [6]
- (ii) Solve the equation fg(x) = 5 when k = 6. [3]
- (iii) Express  $g^{-1}(x)$  in terms of x.

4

[4]

[4]

[2]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.