

## Location Entry Codes

---

As part of CIE's continual commitment to maintaining best practice in assessment, CIE uses different variants of some question papers for our most popular assessments with large and widespread candidature. The question papers are closely related and the relationships between them have been thoroughly established using our assessment expertise. All versions of the paper give assessment of equal standard.

The content assessed by the examination papers and the type of questions is unchanged.

This change means that for this component there are now two variant Question Papers, Mark Schemes and Principal Examiner's Reports where previously there was only one. For any individual country, it is intended that only one variant is used. This document contains both variants which will give all Centres access to even more past examination material than is usually the case.

The diagram shows the relationship between the Question Papers, Mark Schemes and Principal Examiners' Reports that are available.

Question Paper	Mark Scheme	Principal Examiner's Report
Introduction	Introduction	Introduction
First variant Question Paper	First variant Mark Scheme	First variant Principal Examiner's Report
Second variant Question Paper	Second variant Mark Scheme	Second variant Principal Examiner's Report

### Who can I contact for further information on these changes?

Please direct any questions about this to CIE's Customer Services team at:

[international@cie.org.uk](mailto:international@cie.org.uk)

The titles for the variant items should correspond with the table above, so that at the top of the first page of the relevant part of the document and on the header, it has the words:

- First variant Question Paper / Mark Scheme / Principal Examiner's Report

or

- Second variant Question Paper / Mark Scheme / Principal Examiner's Report

as appropriate.



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
International General Certificate of Secondary Education

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 2 9 4 2 3 9 7 2 8 8 \*

**CHEMISTRY**

**0620/31**

Paper 3 (Extended)

**October/November 2008**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES**

Answer **all** questions.

A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part questions.

For Examiner's Use	
<b>1</b>	
<b>2</b>	
<b>3</b>	
<b>4</b>	
<b>5</b>	
<b>6</b>	
<b>7</b>	
<b>Total</b>	

This document consists of **12** printed pages.



1 Complete the following table.

gas	test for gas
ammonia	
	bleaches damp litmus paper
hydrogen	
	relights a glowing splint
	turns limewater milky

[Total: 5]

For  
Examiner's  
Use

2 There are three types of giant structure – ionic, metallic and macromolecular.

(a) Sodium nitride is an ionic compound. Draw a diagram that shows the formula of the compound, the charges on the ions and the arrangement of the valency electrons around the negative ion.

Use x to represent an electron from a sodium atom.  
Use o to represent an electron from a nitrogen atom.

[3]

(b) (i) Describe metallic bonding.

.....  
..... [3]

(ii) Use the above ideas to explain why  
metals are good conductors of electricity,

..... [1]  
metals are malleable.

..... [2]

(c) Silicon(IV) oxide has a macromolecular structure.

(i) Describe the structure of silicon(IV) oxide (a diagram is not acceptable).

.....  
.....  
..... [3]

(ii) Diamond has a similar structure and consequently similar properties.  
Give two physical properties common to both diamond and silicon(IV) oxide.

.....  
..... [2]

[Total: 14]

3 Steel is an alloy made from impure iron.

(a) Both iron and steel rust. The formula for rust is  $\text{Fe}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$ .  
It is hydrated iron(III) oxide.

(i) Name the **two** substances that must be present for rusting to occur.

..... [2]

(ii) Painting and coating with grease are two methods of preventing iron or steel from rusting. Give **two** other methods.

.....  
..... [2]

(b) (i) Name a reagent that can reduce iron(III) oxide to iron.

..... [1]

(ii) Write a symbol equation for the reduction of iron(III) oxide,  $\text{Fe}_2\text{O}_3$ , to iron.

..... [2]

(c) (i) Calculate the mass of one mole of  $\text{Fe}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$ .

..... [1]

(ii) Use your answer to (i) to calculate the percentage of iron in rust.

.....  
..... [2]

(d) Iron from the blast furnace is impure. Two of the impurities are carbon and silicon. These are removed by blowing oxygen through the molten iron and adding calcium oxide.

(i) Explain how the addition of oxygen removes carbon.

.....  
..... [1]


(ii) Explain how the addition of oxygen and calcium oxide removes silicon.

.....  
..... [2]

[Total: 13]

- 4 Across the world, food safety agencies are investigating the presence of minute traces of the toxic hydrocarbon, benzene, in soft drinks. It is formed by the reduction of sodium benzoate by vitamin C.

For  
Examiner's  
Use



- (a) Sodium benzoate is a salt, it has the formula  $C_6H_5COONa$ . It can be made by the neutralisation of benzoic acid by sodium hydroxide.

(i) Deduce the formula of benzoic acid.

..... [1]

(ii) Write a word equation for the reaction between benzoic acid and sodium hydroxide.

..... [1]

(iii) Name **two** other compounds that would react with benzoic acid to form sodium benzoate.

..... [2]

- (b) Benzene contains 92.3% of carbon and its relative molecular mass is 78.

(i) What is the percentage of hydrogen in benzene?

..... [1]

(ii) Calculate the ratio of moles of C atoms: moles of H atoms in benzene.


.....  
..... [2]

(iii) Calculate its empirical formula and **then** its molecular formula.

The empirical formula of benzene is .....

The molecular formula of benzene is ..... [2]

(c) The structural formula of Vitamin C is drawn below.



For  
Examiner's  
Use

(i) What is its molecular formula?

..... [1]

(ii) Name the two functional groups which are circled.

..... [2]

[Total: 12]

5 The electrolysis of concentrated aqueous sodium chloride produces three commercially important chemicals hydrogen, chlorine and sodium hydroxide.


For  
Examiner's  
Use

(a) The ions present are  $\text{Na}^+(\text{aq})$ ,  $\text{H}^+(\text{aq})$ ,  $\text{Cl}^-(\text{aq})$  and  $\text{OH}^-(\text{aq})$ .

(i) Complete the ionic equation for the reaction at the negative electrode (cathode).



(ii) Complete the ionic equation for the reaction at the positive electrode (anode).



(iii) Explain why the solution changes from sodium chloride to sodium hydroxide.

..... [1]

(b) (i) Why does the water supply industry use chlorine?

..... [1]

(ii) Name an important chemical that is made from hydrogen.

..... [1]

(iii) How is sodium hydroxide used to make soap?


..... [2]

[Total: 7]



6 The reactivity series lists metals in order of reactivity.

- (a) To find out which is the more reactive metal, zinc or tin, the following experiment could be carried out.



This experiment could be carried out with other metals and the results recorded in a table. Then the order of reactivity can be deduced.

- (i) The order was found to be:
- |           |                |
|-----------|----------------|
| manganese | most reactive  |
| zinc      |                |
| tin       |                |
| silver    | least reactive |

Complete the table of results from which this order was determined.

aqueous solution	tin Sn	manganese Mn	silver Ag	zinc Zn
tin(II) nitrate		R	NR	R
manganese(II) nitrate				
silver(I) nitrate				
zinc nitrate				

[3]

- (ii) Write the ionic equation for the reaction between tin atoms and silver(I) ions.

.....

[2]

(iii) The following is a redox reaction.



Indicate on the equation the change which is oxidation.  
Give a reason for your choice.

..... [2]

(iv) Explain why experiments of this type cannot be used to find the position of aluminium in the reactivity series.

.....  
..... [2]

(b) Potassium and calcium are very reactive metals at the top of the series. Because their ions have different charges,  $\text{K}^+$  and  $\text{Ca}^{2+}$ , their compounds behave differently when heated.

(i) Explain why the ions have different charges.

.....  
..... [2]

(ii) Their hydroxides are heated.  
If the compound decomposes, complete the word equation.  
If it does not decompose, write "no reaction".

Potassium hydroxide  $\longrightarrow$  .....

Calcium hydroxide  $\longrightarrow$  ..... [2]

(iii) Complete the equations for the decomposition of their nitrates.

$2\text{KNO}_3 \longrightarrow$  ..... + .....

$2\text{Ca}(\text{NO}_3)_2 \longrightarrow$  ..... + ..... + ..... [4]


[Total: 17]

- 7 The alkanes are generally unreactive. Their reactions include combustion, substitution and cracking.

For  
Examiner's  
Use

(a) The complete combustion of an alkane gives carbon dioxide and water.

- (i) 10 cm<sup>3</sup> of butane is mixed with 100 cm<sup>3</sup> of oxygen, which is an excess. The mixture is ignited. What is the volume of unreacted oxygen left and what is the volume of carbon dioxide formed?



Volume of oxygen left = ..... cm<sup>3</sup>

Volume of carbon dioxide formed = ..... cm<sup>3</sup> [2]

- (ii) Why is the incomplete combustion of any alkane dangerous, particularly in an enclosed space?

.....  
..... [2]

(b) The equation for a substitution reaction of butane is given below.



- (i) Name the organic product.

..... [1]


- (ii) This reaction does not need increased temperature or pressure. What is the essential reaction condition?

..... [1]

- (iii) Write a different equation for a substitution reaction between butane and chlorine.

..... [1]

- (c) Alkenes are more reactive and industrially more useful than alkanes. They are made by cracking alkanes.



- (i) Draw the structural formula of the polymer poly(propene).

[2]

- (ii) Give the structural formula and name of the alcohol formed when but-1-ene reacts with steam.

name .....

[1]

structural formula

[1]

- (iii) Deduce the structural formula of the product formed when propene reacts with hydrogen chloride.

[1]

[Total: 12]

**DATA SHEET**  
**The Periodic Table of the Elements**

		Group																																																																																																																																																																			
		I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII																																																																																																																																																								
		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td style="width: 10%;">1</td> <td style="width: 10%;">H</td> <td colspan="10"></td> <td style="width: 10%;">2</td> <td style="width: 10%;">He</td> </tr> <tr> <td></td> <td>Hydrogen</td> <td colspan="10"></td> <td></td> <td>Helium</td> </tr> </table>												1	H											2	He		Hydrogen												Helium																																																																																																																												
1	H											2	He																																																																																																																																																								
	Hydrogen												Helium																																																																																																																																																								
		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td style="width: 10%;">3</td> <td style="width: 10%;">Li</td> <td style="width: 10%;">4</td> <td style="width: 10%;">Be</td> <td colspan="8"></td> <td style="width: 10%;">9</td> <td style="width: 10%;">B</td> <td style="width: 10%;">10</td> <td style="width: 10%;">C</td> <td style="width: 10%;">11</td> <td style="width: 10%;">N</td> <td style="width: 10%;">12</td> <td style="width: 10%;">O</td> <td style="width: 10%;">13</td> <td style="width: 10%;">F</td> <td style="width: 10%;">14</td> <td style="width: 10%;">Ne</td> </tr> <tr> <td></td> <td>Lithium</td> <td></td> <td>Beryllium</td> <td colspan="8"></td> <td></td> <td>Boron</td> <td></td> <td>Carbon</td> <td></td> <td>Nitrogen</td> <td></td> <td>Oxygen</td> <td></td> <td>Fluorine</td> <td></td> <td>Neon</td> </tr> <tr> <td></td> <td>7</td> <td></td> <td>9</td> <td colspan="8"></td> <td></td> <td>5</td> <td></td> <td>6</td> <td></td> <td>7</td> <td></td> <td>8</td> <td></td> <td>9</td> <td></td> <td>10</td> </tr> </table>												3	Li	4	Be									9	B	10	C	11	N	12	O	13	F	14	Ne		Lithium		Beryllium										Boron		Carbon		Nitrogen		Oxygen		Fluorine		Neon		7		9										5		6		7		8		9		10																																																																																
3	Li	4	Be									9	B	10	C	11	N	12	O	13	F	14	Ne																																																																																																																																														
	Lithium		Beryllium										Boron		Carbon		Nitrogen		Oxygen		Fluorine		Neon																																																																																																																																														
	7		9										5		6		7		8		9		10																																																																																																																																														
		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td style="width: 10%;">11</td> <td style="width: 10%;">Na</td> <td style="width: 10%;">12</td> <td style="width: 10%;">Mg</td> <td colspan="8"></td> <td style="width: 10%;">17</td> <td style="width: 10%;">Al</td> <td style="width: 10%;">18</td> <td style="width: 10%;">Si</td> <td style="width: 10%;">19</td> <td style="width: 10%;">P</td> <td style="width: 10%;">20</td> <td style="width: 10%;">S</td> <td style="width: 10%;">21</td> <td style="width: 10%;">Cl</td> <td style="width: 10%;">22</td> <td style="width: 10%;">Ar</td> </tr> <tr> <td></td> <td>Sodium</td> <td></td> <td>Magnesium</td> <td colspan="8"></td> <td></td> <td>Aluminium</td> <td></td> <td>Silicon</td> <td></td> <td>Phosphorus</td> <td></td> <td>Sulphur</td> <td></td> <td>Chlorine</td> <td></td> <td>Argon</td> </tr> <tr> <td></td> <td>23</td> <td></td> <td>24</td> <td colspan="8"></td> <td></td> <td>27</td> <td></td> <td>28</td> <td></td> <td>31</td> <td></td> <td>32</td> <td></td> <td>35.5</td> <td></td> <td>40</td> </tr> </table>												11	Na	12	Mg									17	Al	18	Si	19	P	20	S	21	Cl	22	Ar		Sodium		Magnesium										Aluminium		Silicon		Phosphorus		Sulphur		Chlorine		Argon		23		24										27		28		31		32		35.5		40																																																																																
11	Na	12	Mg									17	Al	18	Si	19	P	20	S	21	Cl	22	Ar																																																																																																																																														
	Sodium		Magnesium										Aluminium		Silicon		Phosphorus		Sulphur		Chlorine		Argon																																																																																																																																														
	23		24										27		28		31		32		35.5		40																																																																																																																																														
		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td style="width: 10%;">19</td> <td style="width: 10%;">K</td> <td style="width: 10%;">20</td> <td style="width: 10%;">Ca</td> <td colspan="8"></td> <td style="width: 10%;">25</td> <td style="width: 10%;">V</td> <td style="width: 10%;">26</td> <td style="width: 10%;">Cr</td> <td style="width: 10%;">27</td> <td style="width: 10%;">Mn</td> <td style="width: 10%;">28</td> <td style="width: 10%;">Fe</td> <td style="width: 10%;">29</td> <td style="width: 10%;">Co</td> <td style="width: 10%;">30</td> <td style="width: 10%;">Ni</td> <td style="width: 10%;">31</td> <td style="width: 10%;">Cu</td> <td style="width: 10%;">32</td> <td style="width: 10%;">Zn</td> <td style="width: 10%;">33</td> <td style="width: 10%;">Ga</td> <td style="width: 10%;">34</td> <td style="width: 10%;">Ge</td> <td style="width: 10%;">35</td> <td style="width: 10%;">As</td> <td style="width: 10%;">36</td> <td style="width: 10%;">Se</td> <td style="width: 10%;">37</td> <td style="width: 10%;">Rb</td> <td style="width: 10%;">38</td> <td style="width: 10%;">Sr</td> <td style="width: 10%;">39</td> <td style="width: 10%;">Y</td> <td style="width: 10%;">40</td> <td style="width: 10%;">Zr</td> <td style="width: 10%;">41</td> <td style="width: 10%;">Nb</td> <td style="width: 10%;">42</td> <td style="width: 10%;">Mo</td> <td style="width: 10%;">43</td> <td style="width: 10%;">Tc</td> <td style="width: 10%;">44</td> <td style="width: 10%;">Ru</td> <td style="width: 10%;">45</td> <td style="width: 10%;">Rh</td> <td style="width: 10%;">46</td> <td style="width: 10%;">Pd</td> <td style="width: 10%;">47</td> <td style="width: 10%;">Ag</td> <td style="width: 10%;">48</td> <td style="width: 10%;">Cd</td> <td style="width: 10%;">49</td> <td style="width: 10%;">In</td> <td style="width: 10%;">50</td> <td style="width: 10%;">Sn</td> <td style="width: 10%;">51</td> <td style="width: 10%;">Sb</td> <td style="width: 10%;">52</td> <td style="width: 10%;">Te</td> <td style="width: 10%;">53</td> <td style="width: 10%;">I</td> <td style="width: 10%;">54</td> <td style="width: 10%;">Xe</td> </tr> <tr> <td></td> <td>Rubidium</td> <td></td> <td>Strontium</td> <td colspan="8"></td> <td></td> <td>Vanadium</td> <td></td> <td>Chromium</td> <td></td> <td>Manganese</td> <td></td> <td>Iron</td> <td></td> <td>Cobalt</td> <td></td> <td>Nickel</td> <td></td> <td>Copper</td> <td></td> <td>Zinc</td> <td></td> <td>Gallium</td> <td></td> <td>Germanium</td> <td></td> <td>Arsenic</td> <td></td> <td>Selenium</td> <td></td> <td>Bromine</td> <td></td> <td>Krypton</td> </tr> <tr> <td></td> <td>85</td> <td></td> <td>88</td> <td colspan="8"></td> <td></td> <td>51</td> <td></td> <td>52</td> <td></td> <td>55</td> <td></td> <td>56</td> <td></td> <td>59</td> <td></td> <td>59</td> <td></td> <td>64</td> <td></td> <td>65</td> <td></td> <td>70</td> <td></td> <td>73</td> <td></td> <td>75</td> <td></td> <td>79</td> <td></td> <td>80</td> <td></td> <td>84</td> </tr> </table>												19	K	20	Ca									25	V	26	Cr	27	Mn	28	Fe	29	Co	30	Ni	31	Cu	32	Zn	33	Ga	34	Ge	35	As	36	Se	37	Rb	38	Sr	39	Y	40	Zr	41	Nb	42	Mo	43	Tc	44	Ru	45	Rh	46	Pd	47	Ag	48	Cd	49	In	50	Sn	51	Sb	52	Te	53	I	54	Xe		Rubidium		Strontium										Vanadium		Chromium		Manganese		Iron		Cobalt		Nickel		Copper		Zinc		Gallium		Germanium		Arsenic		Selenium		Bromine		Krypton		85		88										51		52		55		56		59		59		64		65		70		73		75		79		80		84
19	K	20	Ca									25	V	26	Cr	27	Mn	28	Fe	29	Co	30	Ni	31	Cu	32	Zn	33	Ga	34	Ge	35	As	36	Se	37	Rb	38	Sr	39	Y	40	Zr	41	Nb	42	Mo	43	Tc	44	Ru	45	Rh	46	Pd	47	Ag	48	Cd	49	In	50	Sn	51	Sb	52	Te	53	I	54	Xe																																																																																														
	Rubidium		Strontium										Vanadium		Chromium		Manganese		Iron		Cobalt		Nickel		Copper		Zinc		Gallium		Germanium		Arsenic		Selenium		Bromine		Krypton																																																																																																																														
	85		88										51		52		55		56		59		59		64		65		70		73		75		79		80		84																																																																																																																														
		<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td style="width: 10%;">55</td> <td style="width: 10%;">Cs</td> <td style="width: 10%;">56</td> <td style="width: 10%;">Ba</td> <td colspan="8"></td> <td style="width: 10%;">63</td> <td style="width: 10%;">Tl</td> <td style="width: 10%;">64</td> <td style="width: 10%;">Pb</td> <td style="width: 10%;">65</td> <td style="width: 10%;">Bi</td> <td style="width: 10%;">66</td> <td style="width: 10%;">Po</td> <td style="width: 10%;">67</td> <td style="width: 10%;">At</td> <td style="width: 10%;">68</td> <td style="width: 10%;">Fr</td> <td style="width: 10%;">87</td> <td style="width: 10%;">Ra</td> <td style="width: 10%;">88</td> <td style="width: 10%;">Ac</td> </tr> <tr> <td></td> <td>Cesium</td> <td></td> <td>Barium</td> <td colspan="8"></td> <td></td> <td>Thallium</td> <td></td> <td>Lead</td> <td></td> <td>Bismuth</td> <td></td> <td>Polonium</td> <td></td> <td>Astatine</td> <td></td> <td>Radium</td> <td></td> <td>Actinium</td> </tr> <tr> <td></td> <td>133</td> <td></td> <td>137</td> <td colspan="8"></td> <td></td> <td>204</td> <td></td> <td>207</td> <td></td> <td>209</td> <td></td> <td>210</td> <td></td> <td>210</td> <td></td> <td>210</td> <td></td> <td>226</td> <td></td> <td>226</td> <td></td> <td>227</td> <td></td> <td>227</td> </tr> </table>												55	Cs	56	Ba									63	Tl	64	Pb	65	Bi	66	Po	67	At	68	Fr	87	Ra	88	Ac		Cesium		Barium										Thallium		Lead		Bismuth		Polonium		Astatine		Radium		Actinium		133		137										204		207		209		210		210		210		226		226		227		227																																																																		
55	Cs	56	Ba									63	Tl	64	Pb	65	Bi	66	Po	67	At	68	Fr	87	Ra	88	Ac																																																																																																																																										
	Cesium		Barium										Thallium		Lead		Bismuth		Polonium		Astatine		Radium		Actinium																																																																																																																																												
	133		137										204		207		209		210		210		210		226		226		227		227																																																																																																																																						

\*58-71 Lanthanoid series  
†90-103 Actinoid series

a = relative atomic mass  
X = atomic symbol  
b = proton (atomic) number

		140	Ce	141	Pr	142	Nd	143	Pm	144	Sm	145	Eu	146	Gd	147	Tb	148	Dy	149	Ho	150	Er	151	Tm	152	Yb	153	Lu
		58	Cerium	59	Praseodymium	60	Neodymium	61	Promethium	62	Samarium	63	Europlum	64	Gadolinium	65	Terbium	66	Disprosium	67	Holmium	68	Erbium	69	Thulium	70	Ytterbium	71	Lutetium
		90	Th	91	Pa	92	U	93	Np	94	Pu	95	Am	96	Cm	97	Bk	98	Cf	99	Es	100	Fm	101	Md	102	No	103	Lr
		232	Thorium	231	Protactinium	238	Uranium	237	Neptunium	244	Plutonium	243	Americium	250	Curium	259	Berkelium	266	Californium	271	Einsteinium	285	Fermium	289	Mendelevium	290	Nobelium	294	Lawrencium

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
International General Certificate of Secondary Education

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--

\* 7 5 0 7 3 1 6 8 3 0 \*

**CHEMISTRY**

**0620/32**

Paper 3 (Extended)

**October/November 2008**

**1 hour 15 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

**DO NOT WRITE IN ANY BARCODES**

Answer **all** questions.

A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part questions.

For Examiner's Use	
<b>1</b>	
<b>2</b>	
<b>3</b>	
<b>4</b>	
<b>5</b>	
<b>6</b>	
<b>7</b>	
<b>Total</b>	

This document consists of **12** printed pages.



1 Complete the following table.

*For  
Examiner's  
Use*

gas	test for gas
	turns damp red litmus paper blue
	bleaches damp litmus paper
hydrogen	
oxygen	
carbon dioxide	

[Total: 5]

2 There are three types of giant structure – ionic, metallic and macromolecular.

(a) Sodium sulphide is an ionic compound. Draw a diagram that shows the formula of the compound, the charges on the ions and the arrangement of the valency electrons around the negative ion.

Use x to represent an electron from a sodium atom.  
Use o to represent an electron from a sulphur atom.

[3]

(b) (i) Describe metallic bonding.

.....  
..... [3]

(ii) Use the above ideas to explain why  
metals are good conductors of electricity,

..... [1]

metals are malleable.

..... [2]

(c) Silicon(IV) oxide has a macromolecular structure.

(i) **Describe** the structure of silicon(IV) oxide (a diagram is not acceptable).

.....  
.....  
..... [3]

(ii) Diamond has a similar structure and consequently similar properties.  
Give **two** physical properties common to both diamond and silicon(IV) oxide.

.....  
..... [2]

[Total: 14]



3 Steel is an alloy made from impure iron.

(a) Both iron and steel rust. The formula for rust is  $\text{Fe}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$ .  
It is hydrated iron(III) oxide.

(i) Name the **two** substances that must be present for rusting to occur.

..... and ..... [2]

(ii) Painting and coating with grease are two methods of preventing iron or steel from rusting. Give **two** other methods.

.....  
..... [2]

(b) (i) Name a reagent that can reduce iron(III) oxide to iron.

..... [1]

(ii) Write a symbol equation for the reduction of iron(III) oxide,  $\text{Fe}_2\text{O}_3$ , to iron.

..... [2]

(c) (i) Calculate the mass of one mole of  $\text{Fe}_2\text{O}_3 \cdot 2\text{H}_2\text{O}$ .

..... [1]

(ii) Use your answer to (i) to calculate the percentage of water in rust.

.....  
..... [2]

(d) Iron from the blast furnace is impure. Two of the impurities are carbon and silicon. These are removed by blowing oxygen through the molten iron and adding calcium oxide.

(i) Explain how the addition of oxygen removes carbon.

.....  
..... [1]


(ii) Explain how the addition of oxygen and calcium oxide removes silicon.

.....  
..... [2]

[Total: 13]

- 4 Across the world, food safety agencies are investigating the presence of minute traces of the toxic hydrocarbon, benzene, in soft drinks. It is formed by the reduction of sodium benzoate by vitamin C.

For  
Examiner's  
Use



- (a) Sodium benzoate is a salt, it has the formula  $C_6H_5COONa$ . It can be made by the neutralisation of benzoic acid by sodium hydroxide.

- (i) Deduce the formula of benzoic acid.

..... [1]

- (ii) Write a word equation for the reaction between benzoic acid and sodium hydroxide.

..... [1]

- (iii) Name **two** other compounds that would react with benzoic acid to form sodium benzoate.

..... [2]

(b) Benzene contains 92.3% of carbon and its relative molecular mass is 78.

(i) What is the percentage of hydrogen in benzene?

..... [1]

(ii) Calculate the ratio of moles of C atoms: moles of H atoms in benzene.


.....  
..... [2]

(iii) Calculate its empirical formula and **then** its molecular formula.

The empirical formula of benzene is .....

The molecular formula of benzene is ..... [2]

(c) The structural formula of Vitamin C is drawn below.



(i) What is its molecular formula?

..... [1]

(ii) Name the two functional groups which are circled.

..... [2]

[Total: 12]

5 The electrolysis of concentrated aqueous sodium chloride produces three commercially important chemicals; hydrogen, chlorine and sodium hydroxide.

For  
Examiner's  
Use

(a) The ions present are  $\text{Na}^+(\text{aq})$ ,  $\text{H}^+(\text{aq})$ ,  $\text{Cl}^-(\text{aq})$  and  $\text{OH}^-(\text{aq})$ .

(i) Complete the ionic equation for the reaction at the negative electrode (cathode).



(ii) Complete the ionic equation for the reaction at the positive electrode (anode).



(iii) Explain why the solution changes from sodium chloride to sodium hydroxide.

..... [1]

(b) (i) Why does the water supply industry use chlorine?

..... [1]

(ii) Name an important chemical that is made from hydrogen.

..... [1]

(iii) Sodium hydroxide reacts with fats to make soap and glycerine  
What type of compound are fats?

..... [1]


What type of the reaction is this?

..... [1]

[Total : 7]

6 The reactivity series lists metals in order of reactivity.

- (a) To find out which is the more reactive metal, zinc or tin, the following experiment could be carried out.



This experiment could be carried out with other metals and the results recorded in a table. Then the order of reactivity can be deduced.

- (i) The order was found to be:  
 manganese                    most reactive  
 zinc  
 tin  
 silver                         least reactive

Complete the table of results from which this order was determined.

aqueous solution	tin Sn	manganese Mn	silver Ag	zinc Zn
tin(II) nitrate		R	NR	R
manganese(II) nitrate				
silver(I) nitrate				
zinc nitrate				

[3]

- (ii) Write the equation for the reaction between zinc and silver(I) nitrate.

.....

[2]

- (iii) The following is a redox reaction.



Indicate on the equation which reagent is the oxidant or oxidizing agent.  
Give a reason for your choice.

..... [2]

- (iv) Explain why experiments of this type cannot be used to find the position of aluminium in the reactivity series.

.....

..... [2]

- (b) Potassium and calcium are very reactive metals at the top of the series. Because their ions have different charges,  $\text{K}^+$  and  $\text{Ca}^{2+}$ , their compounds behave differently when heated.

- (i) Explain why the ions have different charges.

.....

..... [2]

- (ii) Their hydroxides are heated.  
If the compound decomposes, complete the word equation.  
If it does not decompose, write "no reaction".

Potassium hydroxide  $\longrightarrow$  .....

Calcium hydroxide  $\longrightarrow$  ..... [2]

- (iii) Complete the equations for the decomposition of their nitrates.

$2\text{KNO}_3 \longrightarrow$  ..... + .....

$2\text{Ca}(\text{NO}_3)_2 \longrightarrow$  ..... + ..... + ..... [4]

[Total: 17]

- 7 The alkanes are generally unreactive. Their reactions include combustion, substitution and cracking.

For  
Examiner's  
Use

(a) The complete combustion of an alkane gives carbon dioxide and water.

- (i) 20 cm<sup>3</sup> of butane is mixed with 150 cm<sup>3</sup> of oxygen, which is an excess. The mixture is ignited. What is the volume of unreacted oxygen left and what is the volume of carbon dioxide formed?



Volume of oxygen left = ..... cm<sup>3</sup>

Volume of carbon dioxide formed = ..... cm<sup>3</sup> [2]

- (ii) Why is the incomplete combustion of any alkane dangerous, particularly in an enclosed space?

.....  
..... [2]

(b) The equation for a substitution reaction of butane is given below.



- (i) Name the organic product.

..... [1]


- (ii) This reaction does not need increased temperature or pressure. What is the essential reaction condition?

..... [1]

- (iii) Write a different equation for a substitution reaction between butane and chlorine.

..... [1]

- (c) Alkenes are more reactive and industrially more useful than alkanes. They are made by cracking alkanes.



- (i) Draw the structural formula of the polymer poly(propene).

[2]

- (ii) Give the structural formula and name of the alcohol formed when propene reacts with steam.

name .....

[1]

structural formula

[1]

- (iii) Deduce the structural formula of the product formed when but-1-ene reacts with hydrogen chloride.

[1]

[Total: 12]

For  
Examiner's  
Use



**DATA SHEET**  
**The Periodic Table of the Elements**

		Group														
I	II	III	IV	V	VI	VII	0									
7 <b>Li</b> Lithium 3	9 <b>Be</b> Beryllium 4	1 <b>H</b> Hydrogen 1	11 <b>B</b> Boron 5	12 <b>C</b> Carbon 6	14 <b>N</b> Nitrogen 7	16 <b>O</b> Oxygen 8	19 <b>F</b> Fluorine 9	20 <b>Ne</b> Neon 10								
23 <b>Na</b> Sodium 11	24 <b>Mg</b> Magnesium 12	27 <b>Al</b> Aluminium 13	28 <b>Si</b> Silicon 14	31 <b>P</b> Phosphorus 15	32 <b>S</b> Sulphur 16	35.5 <b>Cl</b> Chlorine 17	40 <b>Ar</b> Argon 18									
39 <b>K</b> Potassium 19	40 <b>Ca</b> Calcium 20	56 <b>Fe</b> Iron 26	55 <b>Mn</b> Manganese 25	59 <b>Co</b> Cobalt 27	59 <b>Ni</b> Nickel 28	64 <b>Cu</b> Copper 29	65 <b>Zn</b> Zinc 30	70 <b>Ga</b> Gallium 31	73 <b>Ge</b> Germanium 32	75 <b>As</b> Arsenic 33	79 <b>Se</b> Selenium 34	80 <b>Br</b> Bromine 35	84 <b>Kr</b> Krypton 36			
85 <b>Rb</b> Rubidium 37	88 <b>Sr</b> Strontium 38	91 <b>Zr</b> Zirconium 40	93 <b>Nb</b> Niobium 41	96 <b>Mo</b> Molybdenum 42	101 <b>Ru</b> Ruthenium 44	106 <b>Pd</b> Palladium 46	108 <b>Ag</b> Silver 47	112 <b>Cd</b> Cadmium 48	115 <b>In</b> Indium 49	119 <b>Sn</b> Tin 50	122 <b>Sb</b> Antimony 51	127 <b>I</b> Iodine 53	131 <b>Xe</b> Xenon 54			
133 <b>Cs</b> Caesium 55	137 <b>Ba</b> Barium 56	178 <b>Hf</b> Hafnium 72	181 <b>Ta</b> Tantalum 73	184 <b>W</b> Tungsten 74	190 <b>Os</b> Osmium 76	195 <b>Pt</b> Platinum 78	197 <b>Au</b> Gold 79	201 <b>Hg</b> Mercury 80	204 <b>Tl</b> Thallium 81	207 <b>Pb</b> Lead 82	209 <b>Bi</b> Bismuth 83	210 <b>Po</b> Polonium 84	210 <b>Rn</b> Radon 86			
226 <b>Ra</b> Radium 88	227 <b>Ac</b> Actinium 89															
*58-71 Lanthanoid series													175 <b>Lu</b> Lutetium 71			
†90-103 Actinoid series													102 <b>No</b> Nobelium 102			
<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td style="width: 20px;">a</td> <td style="width: 20px;"><b>X</b></td> <td style="width: 20px;">b</td> </tr> </table> <p>Key a = relative atomic mass X = atomic symbol b = proton (atomic) number</p>													a	<b>X</b>	b	169 <b>Tm</b> Thulium 69
a	<b>X</b>	b														
													167 <b>Er</b> Erbium 68			
													165 <b>Ho</b> Holmium 67			
													162 <b>Dy</b> Dysprosium 66			
													159 <b>Tb</b> Terbium 65			
													157 <b>Gd</b> Gadolinium 64			
													152 <b>Eu</b> Europium 63			
													150 <b>Sm</b> Samarium 62			
													144 <b>Nd</b> Neodymium 60			
													141 <b>Pr</b> Praseodymium 59			
													140 <b>Ce</b> Cerium 58			
													238 <b>U</b> Uranium 92			
													232 <b>Th</b> Thorium 90			
													94 <b>Pu</b> Plutonium 94			
													95 <b>Am</b> Americium 95			
													96 <b>Cm</b> Curium 96			
													97 <b>Bk</b> Berkelium 97			
													98 <b>Cf</b> Californium 98			
													99 <b>Es</b> Einsteinium 99			
													100 <b>Fm</b> Fermium 100			
													101 <b>Md</b> Mendelevium 101			
													102 <b>No</b> Nobelium 102			

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.