

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.Write in dark blue or black pen.You may use a pencil for any diagrams or graphs.Do not use staples, paper clips, highlighters, glue or correction fluid.DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

If working is needed for any question it must be shown below that question.


Electronic calculators should be used.

If the degree of accuracy is not specified in the question, and if the answer is not exact, give the answer to three significant figures. Give answers in degrees to one decimal place. For π , use either your calculator value or 3.142.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question. The total of the marks for this paper is 70.

This document consists of **12** printed pages.





| | 3 | | | | | |
|---|---|--------------------------|--|--|--|--|
| 4 | $\frac{3}{5}$ | For Examiner's Use | | | | |
| | Which of the following could be a value of <i>p</i> ? | | | | | |
| | $\frac{16}{27}$ 0.67 60% $(0.8)^2$ $\sqrt{\frac{4}{9}}$ | | | | | |
| | Answer[2] | - | | | | |
| 5 | A meal on a boat costs 6 euros (€) or 11.5 Brunei dollars (\$). | | | | | |
| | In which currency does the meal cost less, on a day when the exchange rate is $\pounds 1 = \$1.9037$? Write down all the steps in your working. | | | | | |
| | Answer [2] | | | | | |
| 6 | Use your calculator to find the value of $2^{\sqrt{3}}$. Give your answer correct to 4 significant figures. | - | | | | |
| | Answer[2] | - | | | | |
| | | | | | | |


| 7 | Solve the equation $4x + 6 \times 10^3 = 8 \times 10^4$. Give your answer in standard form. | For Examiner's Use |
|---|--|--------------------------|
| | $Answer x = \qquad [3]$ | -] |
| 8 | p varies directly as the square root of q . p = 8 when $q = 25$. Find p when $q = 100$. | |
| | $Answer p = \qquad [3]$ | ·] |
| 9 | Ashraf takes 1500 steps to walk <i>d</i> metres from his home to the station. Each step is 90 centimetres correct to the nearest 10 cm. Find the lower bound and the upper bound for <i>d</i> . | |
| | Answer $\leq d <$ [3] | -] |

| | Mon | Tue | Wed | Thu | Fri | Sat | Sun |
|--|-------------------------|-------------------------|----------------------|---------|-------------|--------------|------|
| Opening time | 0600 | 0600 | 0600 | 0600 | 0600 | (<i>a</i>) | 0800 |
| Closing time | 2200 | 2200 | 2200 | 2200 | 2200 | 2200 | 1300 |
| (a) The café is open for a total of 100 hours each week. Work out the opening time on Saturday. | | | | | | | |
| (b) The owner decides to close the café at a later time on Sunday. This increases the total number of hours the café is open by 4%. Work out the new closing time on Sunday. | | | | | | | |
| | | | | Answe | r(b) | | |
| | | | | 1115 WC | <i>r(b)</i> | | |
| Rearrange the for | mula $c = -\frac{1}{c}$ | $\frac{4}{a-b}$ to make | te <i>a</i> the subj | | | | |
| Rearrange the for | mula $c = -\frac{1}{c}$ | $\frac{4}{a-b}$ to make | te <i>a</i> the subj | | r(0) | | |

5

For Examiner's Use

[Turn over




© UCLES 2011

0580/22/M/J/11


For

Use

14 60 students recorded their favourite drink.




0580/22/M/J/11



| 19 | f(x) = x^2 g(x) = 2^x h(x) = $2x - 3$ (a) Find g(3). | For Examiner's Use |
|----|---|--------------------------|
| | Answer(a) | |
| | <i>Answer(b)</i> | |
| | <i>Answer(c)</i> [2] | |







| (i) | the bisector of angle ABC, | [2] |
|-----|----------------------------|-----|
|-----|----------------------------|-----|

- (ii) the locus of points which are equidistant from A and from B. [2]
- (b) Shade the region inside the triangle which is nearer to A than to B and nearer to AB than to BC. [1]

Question 21 is printed on the next page.

 $\mathbf{B} = \begin{pmatrix} 6 \\ -4 \end{pmatrix}$

21 (a)

(i) Work out **AB**.

(ii) Work out **BA**.

Answer(a)(ii)

Answer(a)(i)

[2]

[2]

Find \mathbf{C}^{-1} , the inverse of \mathbf{C} .

 $(b) \quad \mathbf{C} = \begin{pmatrix} 3 & 1 \\ 1 & 1 \end{pmatrix}$

Answer(b)

[2]

 $\mathbf{A} = \begin{pmatrix} 2 & 3 \end{pmatrix}$

Examiner's Use

For

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.