MARK SCHEME for the October/November 2008 question paper

MMM. Hiremepapers.com

0606 ADDITIONAL MATHEMATICS

0606/02

Paper 2, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2008 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



UNIVERSITY of CAMBRIDGE International Examinations

Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2008	0606	02

Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Accuracy mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
 B2, 1, 0 means that the candidate can earn anything from 0 to 2.

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2008	0606	02

The following abbreviations may be used in a mark scheme or used on the scripts:

AG	Answer Given on the question paper (so extra checking is needed to ensure that
	the detailed working leading to the result is valid)

- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)

Penalties

- MR -1 A penalty of MR -1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures this is regarded as an error in accuracy.
- OW -1,2 This is deducted from A or B marks when essential working is omitted.
- PA -1 This is deducted from A or B marks in the case of premature approximation.
- S -1 Occasionally used for persistent slackness usually discussed at a meeting.
- EX -1 Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

Page 4		Mark Scheme	Syllabus	Paper	
		IGCSE – October/November 2008		0606	02
1	\mathbf{A}^{-1}	$= \frac{1}{10} \begin{pmatrix} 4 & -6 \\ -7 & 13 \end{pmatrix}$	B1	+B1	
	eva	luate $\mathbf{A}^{-1} \begin{pmatrix} 41\\ 24 \end{pmatrix}$	M	l	
	x =	2, y = 2.5	A1		
					[4]
2	k(2	$(2x-9)^2$	M	l	
	6(2	$(2x-9)^2$	A1		
	sub	stitute $x = 7$ and $\frac{dx}{dt} = 4$ into $\frac{dy}{dt} = \frac{dy}{dt} \times \frac{dx}{dt}$	M	l	
	600	dt dt dx dt	A1		
					[4]
3	elin	hinate y $b^2 + 4ac$	M1	[/[1	
	m^2	b - 4dc or $b + 4dc+10m - 39 * 0 or (5 + m)^2 * 64$	A1	/11	
	fact	orise 3 term quadratic in m or take square root	M	1	
	-13	<i>B</i> < <i>m</i> < 3	A1		[5]
					[5]
	d	1			
4	(i) $\frac{d}{dx}$	$(\ln x) = \frac{1}{x}$	B1		
	1+1	ln x	B1		
	(ii) ∫(1	$(+\ln x)dx = x\ln x(+c)$	M	l	
	∫ln	$x \mathrm{d}x = x \ln x - \int 1 \mathrm{d}x(+c)$	M	l	
	$x \ln x$	x - x(+c)	A1		
					[5]
5	(i) exp	ress as powers of 2 (or 4 or 8)	M		
	app 7	lies rules of indices $\left[2x - (5 - x) = 4x - 3(x - 3)\right]$		A1	
	1		AI		
	(ii) lg($2y+10$ + lg y = lg { $y(2y+10)$ } or 2 = lg 100	B1		
	$2y^2$	$x^{2} + 10y = 100$ oe	B1		
	5 01	пу	BI		[6]

Page 5		e 5	Mark Scheme	Syllabus	Paper
			IGCSE – October/November 2008	0606	02
6	(a)	10, 3 a multir 450	and 15 B bly 3 values M A	1 [1 1	
	(b)	4×(5 240	×4×3) B B	1+B1 1	[6]
7	(i)	speed <u>1.4</u>	of travel = 4.8 or distance downstream = 14 B OR (4.8) (4.8) (4.8) (4.8) 1.4	1	
	(ii)	$\frac{\text{draw n}}{\sqrt{1.4^2}}$ $\tan^{-1} \cdot \frac{1}{\sqrt{3.7}}$	right angle triangle with 1.4 and (4.8) at 90°B $+(4.8)^2$ M (4.8) A (4.8) 0e 1.4 0er 1.29 radiansA	1 11 11	[6]
8	(i)	5	В	1	
	(ii)	180 о	Еπ В	1	
	(iii)	8 and	-2 B	1+B1	
	corree 2 cyc corree	ct start a les in 0 ct max a	nd endpointsBto 2π Bnd min pointsB	1 1 1	[7]

Page 6		6	Mark Scheme	Syllabus	Paper	
	Ŭ		IGCSE – October/November 2008	0606	02	
_						
9		elimir	hate y (or x)	M1		
		$7x^2 - $	$42x + 35 = 0$ (or $7y^2 + 42y - 49 = 0$)oe	A1		
		solve	3 term quadratic	M1		
		x = 1	and 5 (or $y = -7$ and 1)	A1		
		find s	econd coordinates	M1		
		find n	nid-point	M1		
		use m	m_{AB} , $m_1 m_2 = -1$ and coordinates of a point	Ml		
		<i>y</i> +3	$= -\frac{1}{2}(x-3) \text{ or } x+2y+3 = 0 \text{ or } y = -\frac{1}{2}x-\frac{3}{2}$	A1		
					[8]	
4.0		dy		D.I.		
10	(i)	$\frac{y}{\mathrm{d}x} =$	$3x^2 - 16x + 16$	BI		
		equate	e to 0 and solve 3 term quadratic	M1		
		x = 4	, y = 0	A1 AG		
		4	13 = 256 = 0.48 = 0.5	A 1		
		$x = \frac{1}{3}$	$y = 9\frac{1}{27}$ or $\frac{1}{27}$ or 9.48 or 9.5	AI		
	(ii)	integr	rate	M1		
	()	x^4	$8x^3$			
		4	$\frac{3}{3} + 8x^2$	Al		
		use li	mits of 4 (and 0)	DM1		
		$21\frac{1}{2}$	or 21.3	A1		
		3			[8]	
					[0]	
11	(i)	nlot r	wagainst 1/r with linear scales	M1		
	(1)	xv	4.5 3.24 2.82 2.64			
		1/x	0.5 0.25 0.17 0.125 A	A2, 1, 0		
	(ii)	attem	nt at gradient using plotted points	DM1		
	(11)	5+0	2	A1		
		interc	ent $2+0.1$	B1		
		(or A	1 if calculated from $v = mx + c$)			
		use Y	T = mX + c in correct way	M1		
		450 1	$5 \ 2 \ 5+2x \ 1(5)$			
		$y = \frac{1}{x}$	$\frac{1}{x^2} + \frac{1}{x}$ or $y = \frac{1}{x^2}$ or $y = \frac{1}{x} \left(\frac{1}{x} + 2 \right)$	A1√		
	(iii)	read f	from graph or substitute in formula to find r	M1		
	()	x = 2	$.5\pm0.2$	Al		
		y = 1	$.6 \pm 0.1$	A1		
		-			[11]	

Page 7		Mark Scheme		Syllabus	Paper
		IGCSE – October/November 2008		0606	02
2	EITH	IER			
(i)	$\frac{OC}{2} =$	$= \cos 0.6 \text{ or } OC = 2\cos 0.6 \text{ or } \frac{OC}{\sin 0.97} = \frac{2}{\sin \frac{\pi}{2}}$	M1		
	1.65	2	A1		
	CD =	$2\sin 0.6$ or $CD = \sqrt{OD^2 - OC^2}$	M1		
	1.13		A1		
(ii)	6×0.0	6	B1		
	comp	lete plan $CD + 4 + r\theta + (6 - 1.65)$	M1		
	13.1		A1		
(iii)	$\frac{1}{2} \times 6^2$	$^2 \times 0.6$	B1		
	comp	lete plan $\frac{1}{2}r^2\theta - \frac{1}{2} \times OC \times CD$	M1		
	9.87	2 2	A1		
					[10

OR

(i)	$2t^2 - 12t + 16$ equate to 0 and solve quadratic for 2 values 2 and 4	B1+B1+B1 M1 A1	
(ii)	$s = \int v \mathrm{d}t$	M1	
	$\frac{2}{3}t^3 - 6t^2 + 16t$	A 2, 1, 0√	
	use limits and subtract	DM1	
	$2\frac{2}{2}$ or 2.67	A1	
	3		54.03

[10]