

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

**MARK SCHEME for the October/November 2010 question paper
for the guidance of teachers**

0606 ADDITIONAL MATHEMATICS

0606/13

Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0606	13

Mark Scheme Notes

Marks are of the following three types:

M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

B Accuracy mark for a correct result or statement independent of method marks.

- When a part of a question has two or more “method” steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol \surd implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously “correct” answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
B2, 1, 0 means that the candidate can earn anything from 0 to 2.

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0606	13

The following abbreviations may be used in a mark scheme or used on the scripts:

AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no “follow through” from a previous error is allowed)
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)

Penalties

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become “follow through $\sqrt{\quad}$ ” marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy.
OW –1,2	This is deducted from A or B marks when essential working is omitted.
PA –1	This is deducted from A or B marks in the case of premature approximation.
S –1	Occasionally used for persistent slackness – usually discussed at a meeting.
EX –1	Applied to A or B marks when extra solutions are offered to a particular equation. Again, this is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0606	13

<p>1 $\sec x - \cos x = \frac{1}{\cos x} - \cos x$ $= \frac{1 - \cos^2 x}{\cos x} = \sin x \frac{\sin x}{\cos x}$ $= \sin x \tan x$</p> <p>(Alt: $\frac{\sec^2 x - 1}{\sec x} = \frac{\tan^2 x}{\sec x} = \frac{\sin x}{\cos x} \tan x \cos x$)</p>	<p>M1 M1 A1 [3] M1 M1 A1</p>	<p>M1 for dealing with sec and fractions M1 for use of trig identity M1 for dealing with sec and fractions M1 for use of trig identity</p>
<p>2 (i) ${}^7P_4 = 840$</p> <p>(ii) $4 \times {}^6P_3$ or $\frac{4}{7} \times 840$ 480</p>	<p>B1, B1 [2] M1 A1 [2]</p>	<p>B1 for 7P_4 only M1 for a valid method</p>
<p>3 $mx + 2 = x^2 + 12x + 18$ $x^2 + (12 - m)x + 16 = 0$ $(12 - m)^2 = 4 \times 16$ leading to $m = 4, 20$</p> <p>Alt scheme: $m = 2x + 12$ $(2x + 12)x + 2 = x^2 + 12x + 18$ $x = \pm 4$ so $m = 4, 20$</p>	<p>M1 M1 M1, A1 [4] M1 M1 M1 A1 [4]</p>	<p>M1 for equation in x only, allow unsimplified M1 for use of '$b^2 - 4ac$' M1 for solution of quadratic M1 for equating gradients M1 for elimination of m M1 for x and subsequent calculation for m</p>
<p>4 $f(2) = 8 + 4k - 10 - 3$ $f(-1) = -1 + k + 5 - 3$ $(4k - 5) = 5(k + 1)$ leading to $k = -10$</p>	<p>M1 M1 M1 A1 [4]</p>	<p>M1 for use of $x = 2$ M1 for use of $x = -1$ M1 for attempt to link the two remainders</p>
<p>5 $a = b^2, 2a - b = 3$ $2b^2 - b - 3 = 0$ or $4a^2 - 13a + 9 = 0$ leading to $a = \frac{9}{4}, b = \frac{3}{2}$</p>	<p>B1, B1 M1 A1, A1 [5]</p>	<p>M1 for solution of equations leading to a quadratic. Final A1 – correct pair only.</p>

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0606	13

<p>6 $x = 2$ or -4 or $-\frac{1}{3}$</p> <p>Either $(x - 2)(3x^2 + 13x + 4)$ or $(x + 4)(3x^2 - 5x - 2)$ or $(3x + 1)(x^2 + 2x - 8)$ $(x - 2)(x + 4)(3x + 1)$</p> <p>$x = 2, -4, -\frac{1}{3}$</p>	<p>B1 M1 A1 M1, A1 A1 [6]</p>	<p>B1 for spotting a solution M1 for attempt to get quadratic factor A1 for correct quadratic factor M1 for dealing with quadratic factor A1 for correct factors A1 for all solutions</p>
<p>7 (i) Graph of modulus function</p> <p>(ii) Straight line graph</p> <p>(iii) $8x = \pm(3x - 5)$ leading to $x = \frac{5}{11}$ or 0.455 only</p>	<p>B1 B1 B1 [3] B1 [1] M1 M1, A1 [3]</p>	<p>B1 for shape B1 for 5 marked on y axis B1 for $\frac{5}{3}$ marked on x axis B1 for straight line with greater gradient M1 for attempt to deal with modulus M1 for solution</p>
<p>8 (a) (i) $f_{\min} = -10$, occurs when $x = -2$</p> <p>(ii) e.g. $x \geq -2$</p> <p>(b) (i) $x = \left(\frac{y}{2} - 1\right)$, leading to $g^{-1}(x) = 2(x + 1)$</p> <p>(ii) $\frac{x^2 - x}{2} - 1 = 2(x + 1)$ leading to $x^2 - 5x - 6 = 0$ solution $x = 6$ and -1</p>	<p>B1 B1 [2] B1 [1] M1 A1 [2] M1 DM1 A1 [3]</p>	<p>Allow any suitable domain that makes f a 1:1 function M1 for a valid method of finding the inverse function M1 for correct order DM1 for solution of quadratic</p>

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0606	13

<p>9 (a) $\int x^{\frac{2}{3}} - 6x^{\frac{1}{3}} + 9 dx = \frac{3}{5}x^{\frac{5}{3}} - \frac{9}{2}x^{\frac{4}{3}} + 9x(+c)$</p> <p>(b) (i) $\frac{dy}{dx} = \sqrt{x^2 + 6} + x \left(\frac{2x}{2\sqrt{x^2 + 6}} \right)$</p> <p>(ii) $\int \frac{x^2 + 3}{\sqrt{x^2 + 6}} dx = \frac{1}{2}x\sqrt{x^2 + 6}$</p>	<p>M1 A2,1,0 [3]</p> <p>M1 A2,1,0 [3]</p> <p>M1 A1 [2]</p>	<p>M1 for expansion and attempt to integrate -1 each error</p> <p>M1 for attempt to differentiate a product. -1 each error</p> <p>M1 for use of their answer to (i)</p>
<p>10 (i) $t = \sqrt{e^5 - 1}$ or $t^2 + 1 = e^5$ $t = 12.1$</p> <p>(ii) distance = $\ln 10 - \ln 5$ = $\ln 2$ or 0.693</p> <p>(iii) $v = \frac{2t}{t^2 + 1}, v = 0.8$</p> <p>(iv) $a = \frac{(t^2 + 1)2 - 2t(2t)}{(t^2 + 1)^2}$ When $t = 2, a = -\frac{6}{25}$, or -0.24</p>	<p>B1 B1 [2]</p> <p>M1 A1 [2]</p> <p>M1, A1 [2]</p> <p>M1, A1 A1 [3]</p>	<p>M1 for $s_3 - s_2$</p> <p>M1 for attempt to differentiate</p> <p>M1 for attempt to differentiate a product or quotient A1 all correct, allow unsimplified</p>
<p>11 (i) $\tan x = \frac{4}{3}, x = 53.1^\circ, 233.1^\circ$</p> <p>(ii) $11 \sin y + 1 = 4(1 - \sin^2 y)$ $(4 \sin y - 1)(\sin y + 3) = 0$ $\sin y = \frac{1}{4}, y = 14.5^\circ, 165.5^\circ$</p> <p>(iii) $\cos\left(2z + \frac{\pi}{3}\right) = -\frac{1}{2}$ $2z + \frac{\pi}{3} = \frac{2\pi}{3}, \frac{4\pi}{3}$ so $z = \frac{\pi}{6}, \frac{\pi}{2}$</p>	<p>M1 A1, $\sqrt{A1}$ [3]</p> <p>M1 M1 A1, $\sqrt{A1}$ [4]</p> <p>B1 M1 A1, A1 [4]</p>	<p>M1 for an equation in tan Follow through on their first answer +180°</p> <p>M1 for use of correct identity M1 for dealing with quadratic Follow through on their 14.5</p> <p>M1 for correct order of operations</p>

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0606	13

<p>12 EITHER</p> <p>(i) $3 = A \sin \frac{\pi}{6} + B \cos \frac{\pi}{4}, 3 = \frac{1}{2}A + \frac{1}{\sqrt{2}}B$</p> $\frac{dy}{dx} = 2A \cos 2x - 3B \sin 3x$ $-4 = 2A \cos \frac{2\pi}{3} - 3B \sin \pi$ $A = 4, B = \sqrt{2}$ <p>(ii) $A = \int_0^{\frac{\pi}{3}} 4 \sin 2x + B \cos 3x \, dx$</p> $= \left[-2 \cos 2x + \frac{B}{3} \sin 3x \right]_0^{\frac{\pi}{3}}$ $= \left(-2 \cos \frac{2\pi}{3} + \frac{B}{3} \sin \pi \right) - (-2), = 3$	<p>M1 A1</p> <p>M1</p> <p>A1</p> <p>A1, A1 [6]</p> <p>M1</p> <p>A2,1,0</p> <p>DM1,A1 [5]</p>	<p>M1 for attempt at substitution A1 for correct equation</p> <p>M1 for attempt to differentiate</p> <p>A1 for all correct</p> <p>A1 for each</p> <p>M1 for attempt to integrate</p> <p>-1 each error</p> <p>DM1 for use of limits</p>
<p>12 OR</p> <p>(i) $\frac{dy}{dx} = 8x - 6x^2$</p> <p>Grad at $A = 2$, perp grad = $-\frac{1}{2}$</p> <p>At A, $y = 2$</p> <p>Equation of normal: $y - 2 = -\frac{1}{2}(x - 1)$</p> <p>$C(0, 2.5)$</p> <p>(ii) $B(2,0)$</p> $A = \frac{1}{2}(2.5 + 2) + \int_1^2 4x^2 - 2x^3 \, dx$ $= 2.25 + \left[\frac{4x^3}{3} - \frac{x^4}{2} \right]_1^2$ $= \frac{49}{12} \text{ or } 4.08$	<p>M1</p> <p>M1</p> <p>B1</p> <p>DM1</p> <p>A1 [5]</p> <p>B1</p> <p>M1</p> <p>M1 A1 DM1</p> <p>A1 [6]</p>	<p>M1 for differentiation</p> <p>M1 for use of $m_1 m_2 = -1$</p> <p>B1 for y coordinate</p> <p>DM1 for finding equation of normal</p> <p>A1 answer given</p> <p>B1 for coords of B</p> <p>M1 for area of trapezium</p> <p>M1 for attempt to integrate A1 all integration correct DM1 for correct use of limits</p>