

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

**MARK SCHEME for the October/November 2010 question paper
for the guidance of teachers**

0607 CAMBRIDGE INTERNATIONAL MATHEMATICS

0607/04

Paper 4 (Extended), maximum raw mark 120

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0607	04

1 (a)	5h 21 min seen 5.35 h seen	M1 M1	Subtracting times (321 M1) Converting minutes part to hours (may be later) (340 ÷ 321 M1)
	340 ÷ 5.35 seen 63.551... or $63\frac{59}{107}$	M1 B1 [4]	340 ÷ their time (× 60 M1)
(b)	(i) 54.0 (54.01 – 54.02)	B2 [2]	If B0, M1 for 0.85×63.55 oe
	(ii) 19 18 ft	B3 ft [3]	ft 340 ÷ their (i) changed to hours and minutes added to 13 00 If B0, M1 for 340 ÷ their (i) (6.29...) or $5.35 \div 0.85$ or $321 \div 0.85 \div 60$ M1 (dep) for changing decimal part to minutes
2 (a)	(i) 93 312	B1 [1]	Accept 93 300 or 93 310
	(ii) 9.3312×10^4 ft	B1 ft [1]	ft their (i) B1 for 9.331, 9.33 or 9.3 all $\times 10^4$
	(b) 9.69(0) to 9.691×10^{-3}	B2 [2]	B1 for 0.00969(0) to 0.009691 implied by 9.69^{-03} SC1 for 9.7×10^{-3} or 9.69×10^3
	(c) 4.57 or 4.573....	B1 [1]	
(d) 4.72 or 4.722 to 4.723	B2 [2]	If B0, M1 for $\log 2000 \div \log 5$ or graph clearly sketched showing intersection	
3 (a)	Sketch of U-shaped parabola intersecting x -axis twice or full correct use of formula with $a = 1$, $b = 2$ and $c = -4$ $\left(\frac{-2 \pm \sqrt{20}}{2}\right)$ or correct use of completing the square –3.24, 1.24	M1 A1 A1 [3]	If M1 A0, SC1 for –3.2 or –3.236... and 1.2 or 1.236... If M0, SC2 for –3.24 and 1.24 or SC1 for –3.2 or –3.236... and 1.2 or 1.236...
	(b) $-3.24 \leq x \leq 1.24$ ft	B1 ft B1 ft [2]	ft only if two solutions to part (a) Condone < used and allow in words, if clear
4 (a)	Line joining 5 on each axis approx Horizontal line roughly through 1 Line through origin at more than 45° to x -axis	B1 B1 B1 [3]	All may be freehand
	(b) R in correct region oe	B1dep [1]	dep on B3

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0607	04

5	(a)	5 2 3 2.875 (allow 2.87, 2.88 or 2.9) 4	B1 B1 B1 B1 B1 [5]	
	(b)	2	B2 [2]	If B0, M1 for attempting to find a fraction with denominator 72
	(c) (i)	$\frac{1}{8}$ cao	B2 [2]	If B0, B1 for $\frac{9}{72}$ o.e.
	(ii)	45 ft	B1ft [1]	ft their (i) if answer is integer accept $\frac{45}{360}$ [10]
6	(a)	1.15	B3 [3]	If B0, M1 for $20t + 8(3t - 1)$ and M1 (dep) for this equal to 42.6
	(b) (i)	$\frac{15}{y} + \frac{9}{y+2} = 8$ $15(y+2) + 9y = 8y(y+2)$ or $15y + 30 + 9y = 8y^2 + 16y$ $8y^2 - 8y - 30 = 0$ $\therefore 4y^2 - 4y - 15 = 0$	M2 M1 E1 [4]	Allow M1 for l.h.s. Could still be all over $y(y+2)$ and not expanded or partly or fully expanded Correctly established. Need to see 1 correct line and final answer
	(ii)	$(2y - 5)(2y + 3)$	B2 [2]	Allow SC1 for any other $(2y \pm 5)(2y \pm 3)$
	(iii)	2.5(0) ft	B1ft [1]	ft a positive root from (ii) if the only one from two possible roots. [10]
7	(a)	Real numbers oe	B1 [1]	
	(b)	3, 90	B1 B1 [2]	Allow either way round
	(c) (i)	Stretch Factor 2 x-axis invariant	B1 B1 B1 [3]	Independent
	(ii)	Translation $\begin{pmatrix} -60 \\ 0 \end{pmatrix}$	B1 B1 [2]	Must be translation Independent – Allow description in words [8]
8	(a) (i)	Triangle at (-4, 4), (-1, 4), (-1, 5)	B2 [2]	If B0, SC1 for any translation
	(ii)	Triangle at (-1, -2), (-1, -5), (-2, -5)	B2 [2]	If B0, SC1 if two vertices correct
	(b)	Enlargement, (factor) 2, (centre) (4, 0)	B1 B1 B1 [3]	Each B is independent
	(c)	Translation $\begin{pmatrix} 6 \\ -3 \end{pmatrix}$	B1 B1 [2]	B's independent Must be translation but allow description in words [9]

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0607	04

9	(a) 2, 3, 5, 7, 11, 13, 17, 19 (b) All 8 points correctly placed (c) 3, 11, 17, 19 ft (d) 3 ft (e) B only shaded (i.e. parts in A and C not shaded)	B1 [1] B3 [3] B1 ft [1] B1 ft [1] B1 [1]	B2 for 7 correct and B1 for 6 correct isw extras ft their Venn diagram ft their Venn diagram	[7]
10	(a) (i) One pair of angles equal with reason Second pair of angles equal with reason Angles of triangles equal (ii) 18 (b) (i) 50 (ii) 98 (iii) 5.14 (2...) (iv) 4	R1 [1] R1 [1] R1 [3] B2 [2] B1 [1] B2 [2] B2 [2] B1 [1]	Reasons can only be angles in same segment oe and vertically opposite oe, the second only used once Accept anything suggesting angles same Each R is independent If B0, M1 for 2^2 or 0.5^2 seen If B0, M1 for $180 - (32 + \text{their (i)})$ or for angle QPR = 32 seen or for angle PQY = 58 seen (may be on diagram) If B0, M1 for $\cos 50 = RY \div 8$ oe	[11]
11	(a) 3 points correct 2mm accuracy (b) Negative (c) (i) $y = -0.565x + 58.5$ (ii) 30 or 31 cao	P2 [2] B1 [1] B1 B1 [2] B2 [2]	P1 for 2 correct Allow description e.g. cold goes down as hot goes up Must be in form $mx + c$, allow -0.57 or -0.5652 to -0.5651 for m and 58 or 58.48... for c Must be integer If B0, M1 for using their linear regression equation with $x = 50$	[7]
12	(a) 0.8333 (b) (i) Two accurate points marked C_1 and C_2 (ii) 56.4, 123.6 (iii) 67.2 ft	B3 [3] B1 B1 [2] B1 B1 [2] B1 ft [1]	SC2 for $\frac{5}{6}$, 0.83, 0.833, 0.8333... isw if angle given If B0 and SC0, M1 for $\frac{\sin C}{10} = \frac{\sin 30}{6}$ oe (can be implicit) 2 mm accuracy ft the difference between their answers in (ii)	[8]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – October/November 2010	0607	04

13 (a)	982 (981.7 – 981.9)	B2 [2]	If B0, M1 for $0.5 \times \pi \times 25^2$
(b)	295 000 (294 500 – 294 600) ft	B2 ft [2]	ft their (a) $\times 300$ If B0, M1 for their (a) $\times 300$
(c) (i)	106.3 (106.2 – 106.3)	B3 [3]	Allow 106 If B0, M1 for $\cos = \frac{15}{25}$ oe then M1 dep for $\times 2$
(ii)	299.9 to 300.4 ft	B2 ft [2]	ft their (i) If B0, M1 for $0.5 \times 25^2 \times \sin(\text{their(i)})$ or for $0.5 \times 2 \times 20 \times 15$ oe
(iii)	577.8 to 580 ft	B2 ft [2]	ft their (i) If B0, M1 for their (i) $\div 360 \times \pi \times 25^2$
(iv)	277 – 280.1 ft	B1ft [1]	ft their (iii) – their (ii)
(v)	83.1 to 84.03 ft	B2 ft [2]	ft their (ii) $\times 0.3$ oe If B0, M1 for their (ii) $\times 0.3$ oe
[14]			
14 (a)	One curve reasonable shape, roughly approaching $y = 1$ both ends One max in negative x region One minimum just to right of y -axis or on it	B1 B1 B1 [3]	
(b)	(–5.19, 1.24) (–5.193 to –5.192, 1.238 to 1.239)	B2 [2]	Allow –5.2 and 1.2
(c)	0.161 to 1.24 (0.1614 to 0.1615 and 1.238 to 1.239)	B3 [3]	Allow 0.16 and 1.24 If B0, B1 for top value their y -coord of (b) and M1 (indep) for evidence of finding minimum point
(d)	$y = 1$	B1 [1]	
(e)	–1.62(4...)	B2 [2]	If B0, M1 for line with $c = 1$ and positive gradient added to sketch (may be freehand)
[11]			