

CAMBRIDGE INTERNATIONAL EXAMINATIONS

Cambridge Ordinary Level

MARK SCHEME for the May/June 2015 series

5070 CHEMISTRY

5070/42

Paper 4 (Alternative to Practical), maximum raw mark 60

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.


Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

Page 2	Mark Scheme	Syllabus Paper
	Cambridge O Level – May/June 2015	5071

- 1 (a) (i) silver/silvery/grey (1)
- (ii) $2\text{Mg} + \text{O}_2 \rightarrow 2\text{MgO}$ (1)
- (b) hydrogen/ H_2 (1)
pops in flame/burning splint pops/lighted splint pops (1) [2]
- (c) (i) MgO /magnesium oxide/solid/it disappears/dissolves
or a colourless solution/colourless liquid (is formed) (1) [1]
- (ii) $\text{MgO} + \text{H}_2\text{SO}_4 \rightarrow \text{MgSO}_4 + \text{H}_2\text{O}$ (1) [1]
- [Total: 6]
- 2 (a) (i) $\begin{array}{ccc} 32 & 38 & 44 \\ \underline{(20)} & \underline{(20)} & \underline{(20)} \\ \underline{12} & \underline{18} & \underline{24} \end{array}$ all correct (1)
- $\begin{array}{ccc} 32 & 38 & 44 \\ \underline{(20)} & \underline{(20)} & \underline{(20)} \\ \underline{12} & \underline{18} & \underline{24} \end{array}$ all correct (1) [2]
- (ii) exothermic (1) [1]
- (b) (i) $\begin{array}{ccc} 60/12 = 5 & 13.3/1 = 13.3 & 26.7/16 = 1.67 \\ 3 & : & 8 & : & 1 \end{array}$
Empirical Formula = $\text{C}_3\text{H}_8\text{O}$ (1) **Reject** $\text{C}_3\text{H}_7\text{OH}$
- Molecular formula = $\text{C}_3\text{H}_8\text{O}$ (1) [2]
- (ii) **X** = $\text{C}_2\text{H}_5\text{OH}$ or CH_3OH (1) **Z** = $\text{C}_4\text{H}_9\text{OH}$ or $\text{C}_5\text{H}_{11}\text{OH}$ (1)
- Reasons:** e.g. the more carbon atoms in the molecule/
the more carbon-carbon bonds/bigger M_r (reject A_r)/larger molecules
the more the temperature (rise)/more heat given out or reverse argument/more
exothermic (1) [3]
- (c) (i) propanoic (acid) /propionic (acid)
 $\text{C}_2\text{H}_5\text{COOH}$ / $\text{CH}_3\text{CH}_2\text{COOH}$ / $\text{C}_2\text{H}_5\text{CO}_2\text{H}$ / $\text{CH}_3\text{CH}_2\text{CO}_2\text{H}$
(both name and structure required) (1) [1]
- (ii) (acidified) potassium manganate(VII) or KMnO_4 or potassium
permanganate (1)
purple/pink to colourless/decolourised (1)
OR
(acidified) potassium dichromate or $\text{K}_2\text{Cr}_2\text{O}_7$ (1)
orange to green (1)
(in both cases, award of second mark is conditional on first mark being obtained) [2]

Page 3	Mark Scheme	Syllabus Paper
	Cambridge O Level – May/June 2015	507

(d) propyl propanoate (1)



[2]

[Total: 13]

3 (d) (1)

[Total: 1]

4 (d) (1)

[Total: 1]

5 (c) (1)

[Total: 1]

6 (b) (1)

[Total: 1]

7 (b) (1)

[Total: 1]

8 (a) 16.11 g (1)

[1]

(b) filtration / decant(ation) / centrifugation (1)

[1]

(c) colourless / green to purple / pink (1)

[1]

(d) $\begin{array}{ccc} 32.3 & 39.4 & 47(.0) \\ \underline{6.9} & \underline{13.6} & \underline{21.8} \\ \underline{25.4} & \underline{25.8} & \underline{25.2} \end{array}$ 1 mark for each correct row or column
 to the benefit of the candidate (3)

Mean value = 25.3 (1) cm³

[4]

(e) 0.000506 (1) **OR** ecf titre $\times 0.0200 / 1000$

[1]

(f) 0.00253 (1) **OR** ecf (e) $\times 5$

[1]

(g) (i) 0.0253 (1) **OR** ecf (f) $\times 10$

[1]

(ii) 1.42 (1) g **OR** ecf (g)(i) $\times 56$

[1]

(h) 8.79 (1) **OR** ecf (g)(ii) / (a) $\times 100$

[1]

Page 4	Mark Scheme	Syllabus Paper
	Cambridge O Level – May/June 2015	507

- (i) (i) $(\text{NH}_4)_2\text{SO}_4 : 28/132 \times 100 (1) = 21.2\% (1)$
- (ii) ammonium nitrate/urea/ammonia/ammonium phosphate/potassium nitrate etc. (1)

[Total: 15]

9 (a) transition metal/element (ion or compound) absent (1) [1]

(b) (i) white ppt (1)

(ii) soluble (in excess)/dissolves/(colourless)solution (1) [2]

(c) (i) white ppt **AND** (ii) soluble (in excess)/dissolves/(colourless) solution (1) [1]

- (d) M1 (aq) NaOH/sodium hydroxide/ (1)
M2 Al/aluminium (foil)/Devarda's alloy (1)
M3 warm/heat/boil (1) may appear in observations
M4 ammonia/ NH_3 **OR gas** turns litmus blue (1)

ALLOW

Brown ring test: conc. (1) sulfuric acid/ H_2SO_4 (1) iron(II) sulfate/ FeSO_4 (1) brown ring (1) [4]

[Total: 8]

10 (a) 0.63, 0.73, 0.81, 0.81 (1) [2]
0.76, 0.81, 0.81, 0.81 (1)

(b) $\text{CaCO}_3 + 2\text{HCl} \rightarrow \text{CaCl}_2 + \text{H}_2\text{O} + \text{CO}_2$ (1) [1]

(c) carbon dioxide/gas (evolved which)
escapes (from the apparatus)/leaves (the apparatus)/is lost (from the apparatus)/
removed (from the apparatus)/is released into the air/is liberated to the outside (1) [1]

(d) all points plotted correctly (1)

two smooth curves through the points (within one small square)
one mark for each curve (2) [3]

(e) (i) 0.56 (1)g [1]

(ii) $87.50 - 0.60$ (value from candidates graph to \pm half a small square) = 86.9(0) (1)g [1]

(f) increase rate/increase speed/faster (1)
increased surface area/increased area of contact/more contact
between marble and acid (1) [2]

Page 5	Mark Scheme	Syllabus Paper
	Cambridge O Level – May/June 2015	507

- (g) Answers must be consequential on equation in (b) (unless equation is given as part of answer)

For a 1:2 mole ratio

$$0.036 / 2 = 0.018 \text{ mol CaCO}_3$$

$$0.018 \times 100 = 1.8 \text{ (g) (1)}$$

$$10 - 1.8 = 8.2 \text{ (g) CaCO}_3 \text{ (1)}$$

E.c.f for a 1:1 mole ratio

$$0.036 \times 100 = 3.6 \text{ (g) (1)}$$

$$10 - 3.6 = 6.4 \text{ (g) (1)}$$

[2]

[Total: 13]